

PATENT APPLICATION TRANSMITTAL LETTER

Atty's Docket No.: 09820.114

TO: ASSISTANT COMMISSIONER FOR PATENTS

Enclosed is the patent application of Sandra Austin-Phillips, Richard R. Burgess, Thomas L. German and Thomas Ziegelhoffer for **TRANSGENIC PLANTS AS AN ALTERNATIVE SOURCE OF LIGNOCELLULOSIC-DEGRADING ENZYMES**.

Enclosed are:

- ☒ forty-three (43) pages of text
☒ twelve (12) sheets of drawings
☒ eighteen (18) pages of sequence listing
☒ declaration and associated power of attorney, unsigned
☒ two (2) verified statements to establish small entity status under 37 CFR 1.9 and 1.27, unsigned (Independent Inventor and Non-Profit Organization)
☒ sequence list on diskette
☒ statement that computer readable form and paper copy of sequence listing are identical

CLAIMS AS FILED

For	No. Filed	No. Extra
Basic Fee		
Total Claims	26 - 20 =	6
Indep Claims	5 - 3 =	2
Multiple Dependent Claim Present		

SMALL ENTITY


Rate	Fee
	\$380
x \$9 =	\$
x \$39 =	\$
+ \$130 =	\$
Total	\$

OTHER THAN A
SMALL ENTITY

Rate	Fee
	\$ 760
x \$18 =	\$ 108
x \$78 =	\$ 78
+ \$260 =	\$
Total	\$ 946

- ☒ A check for \$946 to cover the filing fee is enclosed.
- ☒ The Commissioner is hereby authorized to charge payment of the following fees associated with this communication or credit any overpayment to Deposit Account No. 18-2055.
- ☒ Any additional filing fees required under 37 CFR 1.16.
- ☒ Any patent application processing fees under 37 CFR 1.17.

Establishment of a filing date and assignment of a serial number under 37 C.F.R. §1.53 is requested.


 Joseph T. Leone, Reg. No. 37,170
 DEWITT ROSS & STEVENS, S.C.
 Firststar Financial Centre
 8000 Excelsior Drive, Suite 401
 Madison, Wisconsin 53717-1914
 Telephone: (608) 831-2100
 Facsimile: (608) 831-2106

I hereby certify that this correspondence is being deposited with the U.S. Postal Service as Express Mail - Post Office to Addressee, in an envelope addressed to:

Box: Patent Application
 Assistant Commissioner for Patents
 Washington, D.C. 20231.

Express Mail Label No.: EL03623353745Date of Deposit: 8-12-99Signature: Marcia Rayner

08/12/99
 JCS53 U.S. PTO
 09/373272
 08/12/99

Applicant or Patentee: Sandra Austin-Phillips, Richard R. Burgess, Thomas L. German and Thomas Ziegelhoffer

Serial or Patent No.: _____ Atty. Docket No.: 09820.114

Filed or Issued: Simultaneously Herewith

For: **TRANSGENIC PLANTS AS AN ALTERNATIVE SOURCE OF LIGNOCELLULOSIC-DEGRADING ENZYMES**

**VERIFIED STATEMENT (DECLARATION) CLAIMING SMALL ENTITY STATUS
(37 C.F.R. §§1.9(f) and 1.27(d)) - NON-PROFIT ORGANIZATION**

I hereby declare that I am an official empowered to act on behalf of the nonprofit organization identified below:

NAME OF ORGANIZATION Wisconsin Alumni Research Foundation

ADDRESS OF ORGANIZATION P.O. Box 7365, Madison, WI 53707-7365

TYPE OF ORGANIZATION:

- ☐ UNIVERSITY OR OTHER INSTITUTION OF HIGHER EDUCATION
(37 C.F.R. §1.9(e)(1))
- ☒ TAX EXEMPT UNDER IRS CODE (37 C.F.R. 1.9(e)(2))
- ☐ NONPROFIT SCIENTIFIC OR EDUCATIONAL ORGANIZATION UNDER STATUTE
OF STATE OF THE UNITED STATES OF AMERICA (37 C.F.R. (1.9(e)(3))
NAME OF STATE _____
CITATION OF STATUTE _____
- ☐ WOULD QUALIFY AS TAX EXEMPT UNDER IRS CODE IF LOCATED IN THE
UNITED STATES OF AMERICA (37 C.F.R. 1.9(e)(4)/(e)(2))
- ☐ WOULD QUALIFY AS NONPROFIT SCIENTIFIC OR EDUCATIONAL UNDER
STATUTE OR STATE OF THE UNITED STATES OF AMERICA (37 CFR
1.9(e)(4)/(e)(3))
NAME OF STATE _____
CITATION OF STATUTE _____

I hereby declare that the nonprofit organization identified above qualifies as a nonprofit organization as defined in 37 C.F.R. §1.9(e) for purposes of paying reduced fees under 35 USC §41(a) and (b) with respect to the invention entitled **TRANSGENIC PLANTS AS AN ALTERNATIVE SOURCE OF LIGNOCELLULOSIC-DEGRADING ENZYMES** by inventors Sandra Austin-Phillips, Richard R. Burgess, Thomas L. German and Thomas Ziegelhoffer described in the specification filed herewith.

I hereby declare that rights under contract or law have been conveyed to and remain with the nonprofit organization with regard to the above identified invention.

If the rights held by the nonprofit organization are not exclusive, each individual, concern or organization having rights to the invention is listed below and no rights to the invention are held by any person, other than the inventor, who could not qualify as a small business concern under 37 C.F.R. §1.9(d) or by any concern which would not qualify as a small business concern under 37 C.F.R. §1.9(d) or a nonprofit organization under 37 C.F.R. §1.9(e).

FULL NAME N/A

ADDRESS _____

☐ INDIVIDUAL ☐ SMALL BUSINESS CONCERN ☐ NON-PROFIT ORGANIZATION

I acknowledge the duty to file, in this application or patent, notification of any change in status resulting in loss of entitlement to small entity status prior to paying, or at the time of paying, the earliest of the issue fee or any maintenance fee due after the date on which status as a small entity is no longer appropriate. (37 C.F.R. §1.28(b))

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code, and that such willful false statements may jeopardize the validity of the application, any patent issuing thereon, or any patent to which this verified statement is directed.

NAME OF PERSON SIGNING Carl E. Gulbrandsen

TITLE OF PERSON OTHER THAN OWNER Director of Patents and Licensing

ADDRESS OF PERSON SIGNING P.O. Box 7365, Madison, WI 53707-7365

SIGNATURE _____ DATE _____

Applicant or Patentee: Sandra Austin-Phillips, Richard R. Burgess, Thomas L. German and
Thomas Ziegelhoffer

Serial or Patent No.: _____ Atty. Docket No.: 09820.114

Filed or Issued: Simultaneously Herewith

For: **TRANSGENIC PLANTS AS AN ALTERNATIVE SOURCE OF
LIGNOCELLULOSIC-DEGRADING ENZYMES**

**VERIFIED STATEMENT (DECLARATION) CLAIMING SMALL ENTITY STATUS
(37 C.F.R. §§1.9(f) and 1.27(d)) - INDEPENDENT INVENTOR**

As a below named inventor, I hereby declare that I qualify as an independent inventor as defined in 37 C.F.R. §1.9(c) for purposes of paying reduced fees under 35 U.S.C. §41(a) and (b), to the Patent and Trademark Office with regard to the invention entitled **TRANSGENIC PLANTS AS AN ALTERNATIVE SOURCE OF LIGNOCELLULOSIC-DEGRADING ENZYMES** described in the specification filed herewith.

I have not assigned, granted, conveyed or licensed and am under no obligation under contract or law to assign, grant, convey or license any rights in the invention to any person who could not be classified as an independent inventor under 37 C.F.R. §1.9(c) if that person had made the invention, or to any concern which would not qualify as a small business concern under 37 C.F.R. §1.9(d) or a nonprofit organization under 37 C.F.R. §1.9(e).

Each person, concern or organization to which I have assigned, granted, conveyed, or licensed or am under an obligation under contract or law to assign, grant, convey, or license any rights in the invention is listed below.

FULL NAME: Wisconsin Alumni Research Foundation

ADDRESS: P.O. Box 7365, Madison, WI 53717-1914

☐ INDIVIDUAL

☐ SMALL BUSINESS CONCERN

☒ NON-PROFIT ORGANIZATION

I acknowledge the duty to file, in this application or patent, notification of any change in status resulting in loss of entitlement to small entity status prior to paying, or at the time of paying, the earliest of the issue fee or any maintenance fee due after the date on which status as a small entity is no longer appropriate. (37 C.F.R. §1.28(b))

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code, and that such willful false statements may jeopardize the validity of the application, any patent issuing thereon, or any patent to which this verified statement is directed.

<u>Sandra Austin-Phillips</u>	<u>Richard L. Burgess</u>	<u>Thomas L. German</u>
NAME OF INVENTOR	NAME OF INVENTOR	NAME OF INVENTOR

_____ Signature	_____ Signature	_____ Signature
--------------------	--------------------	--------------------

_____ Date	_____ Date	_____ Date
---------------	---------------	---------------

Thomas Ziegelhoffer
NAME OF INVENTOR

Signature

Date

TRANSGENIC PLANTS AS AN ALTERNATIVE SOURCE OF LIGNOCELLULOSIC-DEGRADING ENZYMES

**SANDRA AUSTIN-PHILLIPS
RICHARD R. BURGESS
THOMAS L. GERMAN
THOMAS ZIEGELHOFFER**

This is a continuation-in-part of co-pending application Serial No. 08/883,495, filed June 26, 1997.

This invention was made with United States government support awarded by the following agencies: DOE Grant No. DE-FC05-92OR22072 and USDA Grant Nos. 94-34190-1204 and 92-34190-6941. The United States has certain rights in this invention.

FIELD OF THE INVENTION

The present invention is directed to the production of cellulose-degrading enzymes in genetically recombinant plants and the recombinant plants themselves.

BIBLIOGRAPHY

Complete bibliographic citations for the non-patent references discussed hereinbelow are included in the Bibliography section, immediately preceding the claims. All of the references cited below are incorporated herein by reference.

DESCRIPTION OF THE PRIOR ART

Lignocellulosic plant matter, such as agricultural and forestry waste, as well as energy crops produced specifically for biomass, offer tremendous potential for the renewable production of fuel and as chemical feedstocks. However, production cost for desired products such as alcohols from lignocellulosic material is significantly higher than

the production cost of equivalent alternatives. However, the prospect, either real or perceived, of limited fossil fuel reserves, along with the geo-political issues which swirl about petroleum-producing countries and regions, renders the production of basic chemical feedstocks and fuels from local, renewable sources an attractive alternative to fossil fuels.

For instance, alcohols have the potential to be excellent alternative transportation fuels if their production costs can be lowered. Brazil has sponsored several programs to replace car engines which run on gasoline alone to engines which run on ethanol or a gasoline-ethanol mix.

Unfortunately, the production of ethanol and other feedstock chemicals from lignocellulosic material is far more complex than an analogous production utilizing a starch-based starting material. Compared to lignocellulosic materials, starch is a simple polymer which is readily hydrolyzed to glucose. Yeasts can then be used to convert the glucose to ethanol.

In contrast, lignocellulosic biomass is a much more complex substrate in which crystalline cellulose is embedded within a matrix of hemicellulose and lignin. The intricate structure and relative inaccessibility of these substrates requires pre-treatment for the disruption of the lignocellulosic material, as well as hydrolysis of hemicellulose and lignin into xylose and phenolic compounds, respectively. (See, for instance, *Micelli et al.* (1996), *Belkacemi et al.* (1996), and *Grohmann et al.* (1992).)

Several enzymes which degrade lignocellulosic material, commonly referred to as "cellulases," are known. The term "cellulase" shall be used herein to refer to any and all enzymes which catalyze the cleavage of cellulosic or lignocellulosic materials. Explicitly, but not exclusively, included within this definition are those cellulases which fall under the Enzyme Classification heading EC 3.2.1.x. Various genes encoding cellulases have also been isolated and characterized.

For instance, genes which encode endoglucanases from the fungus *Trichoderma reesei* are known and have been successfully incorporated and expressed in yeast. See, for instance, *Pentilla et al.* (1987). Likewise, cellulase E2 (EC 3.2.1.4) and cellulase

E3 (EC 3.2.1.91) from the thermo-tolerant bacterium *Thermomonospora fusca* are known. See *Lao et al.* (1991), *Spezio et al.* (1993) and *Zhang et al.* (1995).

From a functional viewpoint, cellulases are categorized into two large sub-groups based upon whether they catalyze cleavage from the cellulose chain ends (exocellulases) or if they catalyze cleavage in the middle of the cellulose chain (endocellulases). For instance, cellobiohydrolase I of *T. reesei* (CBH I, EC 3.2.1.91) is an exocellulase, which degrades crystalline cellulose by cleavage from the chain ends. By way of further illustration, CBH I is a 68 kDa protein with a two-domain architecture which is shared by many cellulases. In this chemical architecture, a large catalytic domain is joined to a cellulose-binding domain (CBD) through a flexible linker region. See *Divne et al.* (1994). Similarly, cellulase E3 of *T. fusca* is also an exocellulase.

Different types of cellulases exhibit synergistic activity on complex substrates. This synergism, especially between exocellulases, is believed to be due to differences in their patterns of absorption to and hydrolysis of complex cellulose substrates. See *Henrissat et al.* (1995).

Illustratively, cellulase E2 of *T. fusca* is a 40 kDa endocellulase which cleaves the cellulose chain internally. Such cleavage generates more chain ends for attack by exocellulases. Consequently when CBH I, E2, and E3 cellulases are combined, their activity together is approximately 5-fold greater than their additive individual activities. (See, for instance, *Irwin et al.* (1993) and WO 94/26880.) It is important to note that proteolytic fragments of cellulases can substitute for the intact enzymes in synergistic mixtures. For example, when combined with *T. fusca* E3 and CBH I, the catalytic domain of *T. fusca* E2 ("E2cd") is as active as the intact enzyme in the digestion of filter paper substrate, *Irwin et al.* (1993).

A wide range of compositions containing cellulases are described in the patent literature. For instance, *Evans et al.*, U.S. Patent No. 5,432,074, describe the use of a formulation consisting essentially of a combination of xylanase and xylosidase, but being essentially free of glucanase and cellobiohydrolase. The formulation also contains a lactic acid-producing bacteria. The formulation is used to treat silage to increase its

nutritive value. In operation, the action of the xylanase and xylosidase enzymes degrades non-cellulosic polysaccharides found in the silage material thereby producing sugars for fermentation.

Heterodimers of different types of cellulose-degrading enzymes are described in WO 94/29460. Here, a β -glucosidase molecule and a cellobiohydrolase molecule (*i.e.*, an exocellulase) are chemically bonded to one another by a crosslinking reagent to yield a single molecule which retains the enzymatic activities of the two separate molecules.

Expression constructs which contain cellulase genes for the transformation of yeast have been constructed. For example, *Knowles et al.*, U.S. Patent No. 5,529,919, describe the transformation of *S. cerevisiae* to contain and express a thermostable β -endoglucanase (EG I) of *T. reesei*.

Likewise, attempts have been made to produce transgenic plants which express cellulose-degrading enzymes. *Aspegren et al.* (1995) describe transgenic suspension-cultured barley cells which express EG I of *T. reesei*. The cells were transformed by particle bombardment and transformed cells selected by a co-transformed antibiotic resistance marker. However, no attempt was made to regenerate complete plants from the cultured cells. Of particular note, this reference states that the production of β -glucanases in plant cells may be hampered by the fact that these enzymes catalyze the hydrolysis of essential cell wall components. Attempts by these authors to stably transform tobacco cells with the same construct used to successfully transform the suspended barley cells failed. Here, the authors observed that after transient expression in tobacco protoplasts, cell wall synthesis never resumed.

SUMMARY OF THE INVENTION

The present invention is drawn to genetically recombinant plants which contain one or more exogenous gene sequences which encode one or more cellulose-degrading gene products. The gene product or products are expressed in recoverable quantities in the recombinant plants and can be isolated from the plants, if desired. In the preferred

embodiment, the genetically recombinant plant expresses the gene product constitutively.

However, the invention also encompasses recombinant plants which express the gene product stage-specifically or tissue-specifically. For example, the gene product or products can be expressed in a plant tissue such as the seeds, fruit, leaves, or tubers of the transformed plant host.

The invention is further drawn to recombinant plants as noted above, wherein the plant contains two exogenous genes whose respective gene products are expressed independently of one another. This allows for different types of cellulases to be expressed in different locations within the same recombinant plant. For example, the plant host can be transformed to express two or more heterologous cellulases in different sub-cellular compartments such as the plastid, cytosol, endoplasmic reticulum, mitochondrion, inclusion body, or vacuole. In addition, chloroplast targeting can also be accomplished through the use of direct chloroplast transformation, an approach that circumvents many of the problems associated with expression of heterologous genes in the nuclear genome. *Carrer et al. (1993), McBride et al. (1994).*

The invention is further drawn to a method for producing cellulose-degrading enzymes. The method comprises transforming a plant host with one or more exogenous genes which encode one or more cellulose-degrading gene products such that the gene product or products are expressed in recoverable quantities. The plant matter containing the expressed protein can be used directly as a feedstock for biomass conversion, or, if desired, the exogenous enzymes so produced can be isolated and purified.

The cellulases produced by the transgenic plants of the present invention can be utilized in the same manner as conventionally-derived cellulases. For instance, cellulases produced by the transgenic plants of the present invention can be isolated and used in fermentation processes such as brewing and wine-making. Here, the cellulases function to hydrolyze cellulose and β -glucans during fermentation. Or, as described in Example 4, below, whole plants transformed to express cellulases can be used directly or added to ensiled plant matter to increase the extent of fermentation of the ensiled matter. Plants transformed to express functional cellulases may also be fed directly to livestock, where the cellulase activity aids in the digestion of lignocellulosic substrates.

Cellulases produced in the transgenic plants of the present invention can also be utilized in the production of ethanol and other feedstock chemicals from lignocellulosic substrates.

Cellulases produced by transgenic plants of the present invention can also be used in the textile, pulping, and paper-making industries. For instance, cellulases are conventionally used to treat denim fabrics to give them a "stone-washed" appearance. Cellulases are also used to modify paper pulps by digesting the cellulose fibers contained within the pulp. The cellulases produced by the transgenic plants described herein can be used in this fashion.

BRIEF DESCRIPTION OF THE DRAWINGS

Fig. 1 is a schematic diagram of binary vector T-DNA for an expression construct to transform plants to contain cellulase E2 of *T. fusca*. Promoters and structural genes are depicted as arrows which indicate the direction of transcription. Terminators are depicted as boxes. NPT II = neomycin phosphotransferase; Met-E2m = *T. fusca* E2 (mature form with N-terminal methionine added); MAS-ter = mannopine synthetase terminator; TML-ter = tumor morphology left terminator; MAC-pro = hybrid "MAC" promoter.

Fig. 2 is a schematic diagram of binary vector T-DNA for an expression construct to transform plants to contain cellulase E3 of *T. fusca*. Promoters and structural genes are depicted as arrows which indicate the direction of transcription. Terminators are depicted as boxes. NPT II = neomycin phosphotransferase; Met-E3m = *T. fusca* E3 (mature form with N-terminal methionine added); MAS-ter = mannopine synthetase terminator; TML-ter = tumor morphology left terminator; MAC-pro = hybrid "MAC" promoter.

Fig. 3 is a western blot analysis evidencing the expression of *T. fusca* E2 cellulase in tobacco transformed to contain the expression construct depicted in Fig. 1.

Fig. 4 is a western blot analysis evidencing the expression of *T. fusca* E3 cellulase

in tobacco transformed to contain the expression construct depicted in Fig. 2.

Fig. 5 is a zymogram gel assay evidencing the expression of active *T. fusca* E2 cellulase in alfalfa transformed to contain the expression construct depicted in Fig. 1.

Fig. 6A is a schematic diagram of a binary vector T-DNA for an expression construct to transform plants to contain cellulase E1 of *A. cellulolyticus*.

Fig. 6B is a schematic diagram of a binary vector T-DNA for an expression construct to transform plants to contain the catalytic domain (E1cd) of cellulase E1 of *A. cellulolyticus*.

Fig. 7 is a schematic diagram of binary vector T-DNA for an expression construct to transform plants to contain cellulase CBH I of *T. Reesei*.

Fig. 8A is a plot of data from an activity assay evidencing the expression of *A. cellulolyticus* E1 cellulase in tobacco transformed with Agrobacterium strain PZA8.

Fig. 8B is a plot of data from an activity assay evidencing the expression of *A. cellulolyticus* E1 cellulase in tobacco transformed with Agrobacterium strain PZA9.

Fig. 9 is a western blot analysis evidencing the expression of CBH I cellulase of *T. Reesei* in tobacco transformed to contain the expression construct depicted in Fig. 7.

Fig. 10 is a schematic diagram of binary vector T-DNA for an expression construct to transform plants to contain the CenA endoglucanase of *Cellulomonas fimi*.

Fig. 11 is a schematic diagram of binary vector T-DNA for an expression construct to transform plants to contain endoglucanase D of *Clostridium thermocellum*.

Fig. 12 is a schematic diagram of binary vector T-DNA for an expression construct to transform plants to contain exoglucanase S of *Clostridium cellulovorans*.

Fig. 13 is a schematic diagram of binary vector T-DNA for an expression construct to transform plants to contain exocellulase E6 of *Thermobifida fusca*.

DETAILED DESCRIPTION OF THE INVENTION

The invention is directed to genetically recombinant plants which express one or more exogenous cellulose-degrading (cellulase) enzymes. The invention is further drawn

to a method of producing cellulases in plants. The invention allows the production of cellulases using the means and methods of large-scale agriculture rather than the conventional route of large-scale fermentation of the bacteria or fungi which are native producers of the cellulases.

5 The recombinant plants are produced by incorporating into a plant host genome one or more expression constructs comprising a DNA sequence which encodes a protein having cellulose-degrading activity. Introduction of the exogenous gene or genes into the plant is accomplished by any means known to the art. The expression constructs described hereinbelow enable the stable transformation of plants with one or more genes
10 which encode cellulose-degrading enzymes. The constructs include a DNA coding sequence which encodes a cellulase (as that term is described herein) which is operatively linked to regulatory sequences which direct constitutive, stage-specific, or tissue-specific expression of the cellulase DNA.

Cellulose-Degrading Enzymes (Cellulases) and Genes:

As noted above, the term "cellulase" shall be used herein to refer to any and all enzymes which catalyze the cleavage of cellulosic or lignocellulosic materials. As used herein, "cellulase" is synonymous with "cellulose-degrading enzymes." Explicitly, but not exclusively, included within the term cellulases are those enzymes which fall under the Enzyme Classification heading EC 3.2.1.x. A non-exhaustive list of these enzymes, the genes for all of which can be used in the present invention, includes the following:

Table 1: Polysaccharide-Degrading Enzymes

25	EC 3.2.1.1 (Alpha-amylase)
	EC 3.2.1.2 (Beta-amylase)
	EC 3.2.1.3 (Glucan 1,4-alpha-glucosidase)
	EC 3.2.1.4 (Cellulase, also known as beta-1,4-endoglucanase, <i>e.g.</i> , cellulase E2)
	EC 3.2.1.6 (Endo-1,3(4)-beta-glucanase)
	EC 3.2.1.7 (Inulinase)
30	EC 3.2.1.8 (Endo-1,4-beta-xylanase)
	EC 3.2.1.10 (Oligo-1,6-glucosidase)
	EC 3.2.1.11 (Dextranase)
	EC 3.2.1.14 (Chitinase)

	EC 3.2.1.15 (Polygalacturonase)
	EC 3.2.1.17 (Lysozyme)
	EC 3.2.1.18 (Exo-alpha-sialidase)
5	EC 3.2.1.20 (Alpha-glucosidase)
	EC 3.2.1.21 (Beta-glucosidase)
	EC 3.2.1.22 (Alpha-galactosidase)
	EC 3.2.1.23 (Beta-galactosidase)
	EC 3.2.1.24 (Alpha-mannosidase)
	EC 3.2.1.25 (Beta-mannosidase)
10	EC 3.2.1.26 (Beta-fructofuranosidase)
	EC 3.2.1.28 (Alpha, alpha-trehalase)
	EC 3.2.1.31 (Beta-glucuronidase)
	EC 3.2.1.32 (Xylan endo-1,3-beta-xylosidase)
	EC 3.2.1.33 (Amylo-1,6-glucosidase)
15	EC 3.2.1.35 (Hyaluronoglucosaminidase)
	EC 3.2.1.36 (Hyaluronoglucuronidase)
	EC 3.2.1.37 (Xylan 1,4-beta-xylosidase)
	EC 3.2.1.38 (Beta-D-fucosidase)
	EC 3.2.1.39 (Glucan endo-1,3-beta-D-glucosidase)
20	EC 3.2.1.40 (Alpha-L-rhamnosidase)
	EC 3.2.1.41 (Alpha-dextrin endo-1,6-alpha-glucosidase)
	EC 3.2.1.42 (GDP-glucosidase)
	EC 3.2.1.43 (Beta-L-rhamnosidase)
25	EC 3.2.1.44 (Fucoidanase)
	EC 3.2.1.45 (Glucosylceramidase)
	EC 3.2.1.46 (Galactosylceramidase)
	EC 3.2.1.47 (Galactosylgalactosylglucosylceramidase)
	EC 3.2.1.48 (Sucrose alpha-glucosidase)
	EC 3.2.1.49 (Alpha-N-acetylgalactosaminidase)
30	EC 3.2.1.50 (Alpha-N-acetylglucosaminidase)
	EC 3.2.1.51 (Alpha-L-fucosidase)
	EC 3.2.1.52 (Beta-N-acetylhexosaminidase)
	EC 3.2.1.53 (Beta-N-acetylgalactosaminidase)
	EC 3.2.1.54 (Cyclomaltodextrinase)
35	EC 3.2.1.55 (Alpha-N-arabinofuranosidase)
	EC 3.2.1.56 (Glucuronosyl-disulfoglucosamine glucuronidase)
	EC 3.2.1.57 (Isopullulanase)
	EC 3.2.1.58 (Glucan 1,3-beta-glucosidase)
	EC 3.2.1.59 (Glucan endo-1,3-alpha-glucosidase)
40	EC 3.2.1.60 (Glucan 1,4-alpha-maltotetrahydrolase)
	EC 3.2.1.61 (Mycodextranase)
	EC 3.2.1.62 (Glycosylceramidase)
	EC 3.2.1.63 (1,2-Alpha-L-fucosidase)
	EC 3.2.1.64 (2,6-Beta-fructan 6-levanbiohydrolase)

	EC 3.2.1.65 (Levanase)
	EC 3.2.1.66 (Quercitrinase)
	EC 3.2.1.67 (Galacturan 1,4-alpha-galacturonidase)
	EC 3.2.1.68 (Isoamylase)
5	EC 3.2.1.70 (Glucan 1,6-alpha-glucosidase)
	EC 3.2.1.71 (Glucan endo-1,2-beta-glucosidase)
	EC 3.2.1.72 (Xylan 1,3-beta-xylosidase)
	EC 3.2.1.73 (Licheninase)
	EC 3.2.1.74 (Glucan 1,4-beta-glucosidase)
10	EC 3.2.1.75 (Glucan endo-1,6-beta-glucosidase)
	EC 3.2.1.76 (L-iduronidase)
	EC 3.2.1.77 (Mannan 1,2-(1,3)-alpha-mannosidase)
	EC 3.2.1.78 (Mannan endo-1,4-beta-mannosidase)
	EC 3.2.1.80 (Fructan beta-fructosidase)
15	EC 3.2.1.81 (Agarase)
	EC 3.2.1.82 (Exo-poly-alpha-galacturonosidase)
	EC 3.2.1.83 (Kappa-carrageenase)
	EC 3.2.1.84 (Glucan 1,3-alpha-glucosidase)
	EC 3.2.1.85 (6-Phospho-beta-galactosidase)
20	EC 3.2.1.86 (6-Phospho-beta-glucosidase)
	EC 3.2.1.87 (Capsular-polysaccharide endo-1,3-alpha-galactosidase)
	EC 3.2.1.88 (Beta-L-arabinosidase)
	EC 3.2.1.89 (Arabinogalactan endo-1,4-beta-galactosidase)
	EC 3.2.1.90 (Arabinogalactan endo-1,3-beta-galactosidase)
25	EC 3.2.1.91 (Cellulose 1,4-beta-cellobiosidase, also known as beta-1,4-exocellulases; cellobiohydrolases; and exoglucanases; <i>e.g.</i> , cellulase E3, CBH I)
	EC 3.2.1.92 (Peptidoglycan beta-N-acetylmuramidase)
	EC 3.2.1.93 (Alpha,alpha-phosphotrehalase)
	EC 3.2.1.94 (Glucan 1,6-alpha-isomaltosidase)
30	EC 3.2.1.95 (Dextran 1,6-alpha-isomaltotriosidase)
	EC 3.2.1.96 (Mannosyl-glycoprotein endo-beta-N-acetylglucosamidase)
	EC 3.2.1.97 (Glycopeptide alpha-N-acetylgalactosaminidase)
	EC 3.2.1.98 (Glucan 1,4-alpha-maltohexaosidase)
	EC 3.2.1.99 (Arabinan endo-1,5-alpha-L-arabinosidase)
35	EC 3.2.1.100 (Mannan 1,4-beta-mannobiosidase)
	EC 3.2.1.101 (Mannan endo-1,6-beta-mannosidase)
	EC 3.2.1.102 (Blood-group-substance endo-1,4-beta-galactosidase)
	EC 3.2.1.103 (Keratan-sulfate endo-1,4-beta-galactosidase)
	EC 3.2.1.104 (Steryl-beta-glucosidase)
40	EC 3.2.1.105 (Strictosidin beta-glucosidase)
	EC 3.2.1.106 (Mannosyl-oligosaccharide glucosidase)
	EC 3.2.1.107 (Protein-glucosylgalactosylhydroxylysine glucosidase)
	EC 3.2.1.108 (Lactase)
	EC 3.2.1.109 (Endogalactosaminidase)

EC 3.2.1.111 (Mucinaminylserine mucinaminidase)
 EC 3.2.1.111 (1,3-Alpha-L-fucosidase)
 EC 3.2.1.112 (Deoxglucosidase)
 EC 3.2.1.113 (Mannosyl-oligosaccharide 1,2-alpha-mannosidase)
 EC 3.2.1.114 (Mannosyl-oligosaccharide 1,3-1,6-alpha-mannosidase)
 EC 3.2.1.115 (Branched-dextran exo-1,2-alpha-glucosidase)
 EC 3.2.1.116 (Glucan 1,4-alpha-maltotriohydrolase)
 EC 3.2.1.117 (Amygdalin beta-glucosidase)
 EC 3.2.1.118 (Prunasin beta-glucosidase)
 EC 3.2.1.119 (Vicianin beta-glucosidase)
 EC 3.2.1.120 (Oligoxyloglucan beta-glycosidase)
 EC 3.2.1.121 (Polymannuronate hydrolase)
 EC 3.2.1.122 (Maltose-6'-phosphate glucosidase)
 EC 3.2.1.123 (Endoglycosylceramidase)
 EC 3.2.1.124 (3-Deoxy-2-octulosonidase)
 EC 3.2.1.125 (Raucaffricine beta-glucosidase)
 EC 3.2.1.126 (Coniferin beta-glucosidase)
 EC 3.2.1.122 (1,6-Alpha-L-fucosidase)
 EC 3.2.1.128 (Glycyrrhizinate beta-glucuronidase)
 EC 3.2.1.129 (Endo-alpha-sialidase)
 EC 3.2.1.130 (Glycoprotein endo-alpha-1,2-mannosidase)
 EC 3.2.1.131 (Xylan alpha-1,2-glucuronosidase)
 EC 3.2.1.132 (Chitosanase)
 EC 3.2.1.133 (Glucan 1,4-alpha-maltohydrolase)
 EC 3.2.1.134 (Difructose-anhydride synthase)
 EC 3.2.1.135 (Neopullulanase)
 EC 3.2.1.136 (Glucuronoarabinoxylan endo-1,4-beta-xylanase)
 EC 3.2.1.137 (Mannan exo-1,2-1,6-alpha-mannosidase)
 EC 3.2.1.138 (Anhydrosialidase)

DNA sequences encoding enzymes having any of the above-described functionalities can be obtained from several microbial sources, including bacterial and fungal sources. Cloning the gene or cDNA sequence of the desired enzyme can be achieved by several well-known methods. A preferred method is to purify the cellulase of interest (or purchase a sample if commercially available) and determine its N-terminal amino acid sequence, as well as several internal amino acid sequences, using known methods. Oligonucleotide probes corresponding to the amino acid sequence are then constructed (again using known methods) and used to screen a genomic or cDNA library of the organism from which the cellulase was isolated. Positive hybrids are identified,

characterized using known methods (restriction enzyme analysis, etc.), and cloned by known means to yield DNA fragments containing the coding sequence for the desired cellulase activity. (See, for instance, *Current Protocols in Molecular Biology*, Chapters 5 and 6.)

5 If a partial nucleotide sequence of the cellulase of choice is already known, this information can be used to construct suitable primers to directly clone the corresponding cDNA using the polymerase chain reaction (PCR). (See *Current Protocols in Molecular Biology*, Chapter 15.)

10 Particularly preferred for use in the present invention are those enzymes falling within the classifications EC 3.2.1.4; EC 3.2.1.6; EC 3.2.1.21; and EC 3.2.1.91. The functionality of these particular enzymes is summarized as follows:

EC 3.2.1.4 enzymes (β -1,4-endoglucanases) hydrolyze internal 1,4 glycosidic bonds of the polysaccharide chain, thereby yielding new chain ends at the surface of cellulose crystals.

EC 3.2.1.6 enzymes (β -1,3-endoglucanases) hydrolyze internal 1,3 glycosidic bonds of the polysaccharide chain, which also results in the formation of new chain ends at the surface of cellulose crystals.

EC 3.2.1.21 enzymes (β -glucosidases) hydrolyze cellobiose into glucose, a readily fermentable substrate.

EC 3.2.1.91 enzymes (β -1,4-exocellulases) cleave cellobiosyl residues (cellobiose is a glucose dimer) from the chain ends of cellulose.

Particularly preferred enzymes (and hence particularly preferred genes) for use in the present invention are cellulase E2 and cellulase E3 of *T. fusca* and CBH I of *T. reesei*.

25 **Expression Constructs:**

Once the protein coding sequence (*i.e.*, the cellulase gene) has been identified and isolated, it must be inserted into an appropriate expression construct containing regulatory elements to direct the expression of the gene and to direct secretion of the gene product

or targeting of the gene product to a particular sub-cellular location or organelle. Manipulation of oligonucleotide sequences using restriction endonucleases to cleave DNA molecules into fragments and DNA ligase enzymes to unite compatible fragments into a single DNA molecule with subsequent incorporation into a suitable plasmid, cosmid, or other transformation vector are well-known to the art.

A transcription regulatory sequence must be included in the expression construct in order to direct the transformed plant cells to transcribe the inserted cellulase coding sequence. Transcriptional regulators may be inducible or constitutive. Inducible transcription regulators direct transcription of the downstream coding sequences in a tissue-specific or growth-stage specific manner. Constitutive regulators provide for sustained transcription in all cell tissues. For purposes of the present invention, constructs which provide constitutive expression of the coding sequence are preferred.

It is also preferred that the expression construct contain a transcription initiation sequence from the tumor-inducing plasmid (Ti) of *Agrobacterium*. Several T-DNA transcription initiation sequences are well known and include, without limitation, the octopine synthase, nopaline synthase, and mannopine synthase initiators.

Downstream of the initiation sequence and fused to the coding sequence, the expression construct may be manipulated to contain a leader signal sequence which directs the resulting polypeptide to a particular organelle or targets the expressed product for secretion (or to signal post-transcriptional or post-translational modification of the gene product).

Likewise, the expression construct should also include a termination sequence to signal transcription termination.

To facilitate selection of successfully transformed plants, the expression construct should also include one or more selectable markers. The neomycin phosphotransferase gene (NPT II), a well-characterized and widely employed antibiotic resistance selection marker is preferred. This marker provides resistance to kanamycin. A large number of other markers are known and can be used with equal success (*e.g.*, other antibiotic resistance markers, dihydrofolate reductase, luciferase, β -glucuronidase, and the like).

For example, Figs. 1 and 2 depict schematic representations of suitable expression constructs for transformation of plants. These constructs are intended for use with *Agrobacterium*-mediated transformation using the binary vector approach. However, these same constructs can be coated onto micro-projectiles for transformation by particle bombardment. With the exception of the coding sequence, these two constructs are essentially identical: Fig. 1 is a schematic diagram of binary vector T-DNA for an expression construct to transform plants to contain cellulase E2 of *T. fusca*.

Fig. 2 is a schematic diagram of binary vector T-DNA for an expression construct to transform plants to contain cellulase E3 of *T. fusca*.

In both Fig. 1 and Fig. 2, promoters and structural genes are depicted as arrows which indicate the direction of transcription and terminators are depicted as boxes. See the "Brief Description of the Figures" for a legend to the abbreviations. In the expression constructs depicted in Figs. 1 and 2, the "MAC" hybrid promoter drives the transcription of the recombinant cellulase genes. Both constructs also contain a constitutive NPT II expression cassette to allow for antibiotic resistance selection using kanamycin. The coding sequence of the construct shown in Fig. 1 (Met-E2m) encodes cellulase E2 from *T. fusca*. (See SEQ. ID. NO: 1; ATG start codon at nt's 255-257, TGA stop codon at nt's 1578-80, first codon of mature E2 protein (AAT) at nt's 348-350.) This sequence encodes the mature form of the enzyme with an N-terminal methionine added. In the same fashion, the coding sequence of the construct shown in Fig. 2 (Met-E3m) encodes cellulase E3 from *T. fusca*. (See SEQ. ID. NO: 2; ATG start codon at nt's 575-577, TAA stop codon at nt's 2363-65, first codon of mature E3 protein (GCC) at nt's 689-692.) This sequence also encodes the mature form of the enzyme with an N-terminal methionine added.

Further examples of constructs which drive targetted expression of cellulose-degrading enzymes are provided in the Examples hereinbelow. Specifically included in the Examples are transformations illustrating apoplastic targeting and accumulation of two additional cellulases. The first of these is the endoglucanase E1 of *Acidothermus cellulolyticus* (EC 3.2.1.4, SEQ. ID. NO: 8). Also included is the cellobiohydrolase

CBH I of *T. Reesei* (EC 3.2.1.91, SEQ. ID. NO: 9). In addition, further Examples of plant expression constructs containing cellulase genes encoding both endoglucanases and cellobiohydrolases (exoglucanase, exocellulase) are provided.

5 Transformation of Plants:

Transformation of the plants can be accomplished by any means known to the art, including *Agrobacterium*-mediated transformation, particle bombardment, electroporation, and virus-mediated transformation. The method of transformation is not critical to the functionality of the present invention insofar as the method chosen successfully incorporates the oligonucleotide construct containing the cellulase-encoding region and any accompanying regulatory sequences into the plant host. The nature of the plant host to be transformed has some bearing on the preferred transformation protocol. For dicots, *Agrobacterium*-mediated transformation utilizing protoplasts or leaf disks is most preferred. Although the Examples disclose the use of tobacco and alfalfa as bioreactors for cellulase production, any crop plant, including monocots, can be utilized. Transformation of monocots is typically achieved by particle bombardment of embryogenic cell lines or cultured embryos. See, for instance, *Vasil et al.* (1993) and *Castillo et al.* (1994). Recent developments in "super-binary" vectors, however, also allow for the use of *Agrobacterium*-mediated gene transfer in most of the major cereal crops. See, for instance, *Ishida et al.* (1996). In this case, the explant source is typically immature embryos.

Agrobacterium-mediated transformation of the plant host using explants is preferred for its relative ease, efficiency, and speed as compared to other methods of plant transformation. For example, disks are punched from the leaves of the plant host and cultured in a suitable medium where they are then exposed to *Agrobacterium* containing the expression construct and (preferably) a disarmed tumor-inducing (Ti) plasmid. *Agrobacterium tumefaciens* LBA 4404 is the preferred strain for transformation. The preferred binary vector is the pCGN1578 binary vector (*McBride and Summerfelt* (1990)).

The binary vector transformation method is well known and needs only be briefly described herein. See *Zambryski et al.* (1989) for a complete review. The Ti plasmid of *Agrobacterium* contains virulence genes (*vir*) which encode trans-acting proteins that enable the transfer of a portion of the plasmid (the T-DNA) to a plant cell. The T-DNA portion of the Ti plasmid is flanked by two border regions (the right and left borders) which act as recognition sites for the excision of the T-DNA from the plasmid prior to its transfer to the plant host. Excision of the T-DNA is mediated by the *vir* genes of the Ti plasmid and involves nicking of the right and left borders of the T-DNA, which frees a single-stranded oligonucleotide fragment. This fragment is then mobilized out of the *Agrobacterium* and into the plant host target.

In the binary vector method, the T-DNA with its right and left border regions is cloned into *E. coli* in known fashion, and the wild-type genes normally found between the two border regions is excised. The expression construct encoding the cellulase of interest is inserted between the right and left border regions. This construct is designated the "binary plasmid." Construction of the binary plasmid is accomplished utilizing the well-characterized recombinant genetic methods applicable to *E. coli*. Successful transformants are selected utilizing a co-transformed marker appropriate for *E. coli*.

The binary plasmid is then mobilized back into *Agrobacterium*. This is accomplished by direct transformation procedures well known to those skilled in the art.

The *Agrobacterium* itself, such as the preferred LBA 4404 strain, is genetically manipulated to contain a Ti plasmid (called the helper plasmid) which lacks the T-DNA and the tumor-inducing regions (i.e., the Ti plasmid is "disarmed") but which still encodes the virulence proteins necessary for DNA transfer. By cooperation between the helper plasmid and the binary plasmid, the length of DNA between the two border regions of the binary plasmid is excised and mobilized into the plant host, where it is incorporated into the plant host genome. The binary method derives its name from the fact that the plasmid containing the expression construct to be transferred is maintained within *Agrobacterium* as a distinct and independently replicating vector from the Ti plasmid itself.

Selection of successful transformants is accomplished using the co-transformed selection marker discussed above. If the marker is NPT II, selection is accomplished by treatment with kanamycin.

For the present invention, the most preferred plants for transformation are alfalfa and tobacco. However, any plant species will function with comparable success. Included among the plant species which can be utilized in the present invention are cauliflowers, artichokes, apples, bananas, cherries, cucumbers, grapes, lemons, melons, nuts, oranges, peaches, pears, plums, strawberries, tomatoes, cabbages, endive, leeks, lettuce, spinach, arrowroot, beets, carrots, cassava, turnips, radishes, yams, sweet potatoes, beans, peas, soya, wheat, barley, corn, rice, rapeseed, millet, sunflower, oats, tubers, kohlrabi, potatoes, and the like.

The plants to be transformed are preferably common green field plants, such as the preferred alfalfa and tobacco, as well as soya, corn, and the like. Equally preferred are plant hosts which are grown specifically for "biomass energy," such as switchgrass, poplar, and the like. In this instance, the enzymes would not be recovered from the plants. The plants are then transformed and regenerated into whole plants which express fully-functional, cellulose-degrading enzymes in economically significant quantities.

Alfalfa is one of the most preferred plant species for use in the present invention because alfalfa is a hardy, perennial plant, which grows well with minimal fertilization and irrigation. Alfalfa is also a very prolific plant. In temperate areas such as those found in the midwestern United States, alfalfa will yield three or more harvests per growing season. Methods have also been developed for wet fractionation of the herbage matter to recover value-added products therefrom.

Tobacco is equally preferred for its prolific growth, ease of transformation, and its well-characterized genetics. Both alfalfa and tobacco are widely cultivated throughout the United States and in other parts of the world.

In the most preferred embodiment, alfalfa or tobacco plants are stably transformed to express, constitutively, enzymatically active E2 or E3 cellulases from *T. fusca*. Also preferred are alfalfa or tobacco which express enzymatically active CBH I from *T. reesei* or combinations of E2, E3, and CBH I. The *T. fusca* cellulases are most preferred

because they are native to thermo-tolerant bacteria and are relatively heat stable. This allows isolation of the cellulase from plant material using relatively rigorous heat precipitation without adversely affecting the activity of the cellulase.

5 Stage-Specific and Tissue-Specific Expression of Cellulases:

Because the enzymes to be expressed by the transformed plant hosts hydrolyze components of the plant cell wall, high levels of expression might have a deleterious effect on the plant host. Therefore, targeting of the expressed enzyme to particular sub-cellular compartments may be preferred. Targeting of the expressed enzyme may also
10 be preferred to avoid expression of the enzyme in sub-cellular compartments where proteolytic activity is high. Targeting of the expressed enzyme may also be preferred if the exogenous cellulase activity interferes with the normal cellular metabolism of certain compartments.

For instance, targeting expression to the apoplast allows the exogenous protein to avoid the more active protein-degrading systems of other cellular compartments, such as in plant leaf vacuoles.

Several signal sequences are known and can be utilized in the present invention. For example, signal sequences for targeting to the secretory pathway are known, *Wandelt et al.* (1992), *Bednarek* (1991), *Mason et al.* (1988), as are sequences for targeting to the chloroplast, *Keegstra et al.* (1993), and the mitochondrion, *de Castro Silva Filho et al.* (1996).

For apoplast targeting, the VSP leader is preferred. The VSP leader comprises the amino acid sequence: NH₃-Met-Lys-Leu-Phe-Val-Phe-Phe-Val-Ala-Ala-Val-Val-Leu-Val-Ala-Trp-Pro-Cys-His-Gly-Ala- (SEQ. ID. NO: 3). See *Mason et al.* (1988).

25 Additionally, bacterial secretory sequences found in the wild-type cellulase gene may be removed to afford cytoplasmic expression of the enzyme in the recombinant plant host.

Targeting can be achieved by fusing combinations of mitochondrial and chloroplast targeting signals to the N-terminus of the desired cellulase, as has been

demonstrated for the reporter genes chloramphenicol acetyl transferase and β -glucuronidase, *de Castro Silva Filho et al.* (1996). In some cases, efficient translocation requires the presence of both signal peptides, with the amino terminal peptide being crucial in specifying import into a particular organelle. In addition, vacuole targeting can be achieved by fusing the sequence encoding the N-terminal 146 amino acids of the vacuolar patatin protein between a secretory leader and structural gene for the cellulase, as has been demonstrated for the yeast invertase gene, *Sonneveld et. al.* (1991).

Regeneration of Mature Transgenic Plants:

Transgenic tobacco and alfalfa were produced by *Agrobacterium*-mediated transformation using explants as source material. This is a routine method easily followed by those skilled in the art. The production methods for transgenic tobacco and alfalfa are given as non-limiting illustrations of the practice of the invention.

The transformation procedure for tobacco is essentially the explant method developed by Horsh et al. (1985). Leaf explants are taken from the second and third fully expanded leaves of three-week old *in vitro* shoot cultures of *Nicotiana tabacum* W38 maintained on MS medium, *Murashige and Skoog* (1962). The leaf pieces are cut into 1 cm squares and pre-cultured on MS medium with 2.0 mg/L 6-benzyl-aminopurine (BAP) and 0.1 mg/L alpha-naphthalene acetic acid (NAA) for 24 hours at 25°C with a 16 hour photo period of 70-90 $\mu\text{E m}^{-2}\text{s}^{-1}$. After pre-culture, explants are placed into a suspension of *Agrobacterium* cells. After 30 minutes, leaf explants are blotted on filter paper and placed abaxial-side down on MS medium with 1.0 mg/L BAP and 0.1 mg/L NAA and co-cultivated for four days under the same conditions as given above. Leaf pieces are then rinsed three times in sterile water, blotted on filter paper, and transferred to the media used for co-cultivation but containing 100 mg/L kanamycin and 400 mg/L carbenicillin. Plantlets (typically 2-3) develop 10-14 days later from callus formed along cut leaf edges. If desired, further plantlet formation can be achieved by transfer of explants to fresh medium at two week intervals. Plantlets are excised and rooted on MS media containing 100 mg/L kanamycin and 400 mg/L carbenicillin.

To transform alfalfa, new-growth trifoliate are taken from alfalfa plants (regenerable genotypes, *Bingham et al.* (1975)) maintained in a growth room and sterilized using alcohol and bleach washes (30 seconds in 70% alcohol, 90 seconds in 20% hypochlorite + 0.1% SDS, followed by three rinses in sterile distilled water). Leaf edges are cut on moist filter paper and tissue then placed into liquid SH-II medium. (*Bingham et al., supra.*) When sufficient explants have been taken, the explants are moved to a suspension of *Agrobacterium* cells containing the engineered plasmid. (The *Agrobacterium* suspension is taken from an overnight culture grown in liquid YEP selection medium.) Cell density is adjusted to fall between about 0.6 to about 0.8 at A_{660} . After 30 minutes inoculation, the explants are gently blotted on filter paper and placed on B5H medium, *Brown and Atanassov* (1985), for four days. They are then rinsed twice in sterile water and cultured on B5H for a further four days. At the end of this period, they are rinsed three times and transferred to B5H containing 25 mg/L kanamycin and 250 mg/L carbenicillin. Plates are maintained at 24°C, 16 hour photo period, light intensity 60-80 $\mu\text{E m}^{-2}\text{s}^{-1}$. Explant-derived calli (and occasionally embryoids) which form within 3 weeks on this medium are moved to B5H with antibiotics but without hormones to allow for further embryoid production and development of existing embryoids. After three to four weeks, embryos are transferred to MS medium including the two antibiotics to allow for development into plantlets. Callus forms on untreated explants in the presence of 25 mg/L kanamycin but embryos are never produced. Each explant piece can give rise to multiple (up to 40) embryos. Plantlets are rooted on MS medium lacking antibiotics.

Monitoring Cellulase Expression:

Cellulase expression can be monitored using a number of different methods, the two most common being western blot analysis (which detects cellulase protein using antibodies specific for the cellulase of interest) and zymographic analysis or enzyme assay (both of which measure the ability of the expressed cellulase to degrade a cellulosic substrate).

Briefly, in the western blot technique, whole plant samples (or root tips, leaves, etc.) are ground in an extraction buffer (preferably 50 mM sodium acetate (pH 5.5) and 10 mM dithiothreitol) and an aliquot of the extract loaded onto an electrophoresis gel (*e.g.*, polyacrylamide containing SDS). Preferably, identical extractions are performed on non-transformed plants and aliquots of these extractions are then loaded onto parallel lanes of the gel to act as negative controls. Serial dilutions of purified cellulase standards can be also electrophoresed to act as positive controls. The gel is then subjected to electrophoresis in standard and well known fashion.

After electrophoresis is complete, the separated proteins are electro-transferred to a nitrocellulose, PVDF, or nylon membrane, in well known fashion. The membrane containing the immobilized proteins is then immersed in a non-specific blocking buffer or detergent (*e.g.*, "TWEEN 20"), and then placed in a solution containing an antibody (the primary antibody) which is specifically reactive with the particular cellulase under investigation. The membrane is then washed and exposed to an enzyme-antibody conjugate directed against the primary antibody (*e.g.*, goat anti-rabbit IgG). The membrane is then exposed to a chromogenic or luminescent substrate to visualize cellulase hybridization on the membrane.

Zymograms in which the cellulase of interest is resolved in a gel system and then assayed for activity within the gel provide a relatively simple way to assess the activity of cellulases in crude cell lysates. See *Coughlan* (1988). In this approach, plant tissue is ground in the presence of an appropriate grinding buffer (100 mM Tris-HCl pH 9.0, 5 mM 2-mercaptoethanol, 1 mM phenylmethanesulfonyl fluoride, 0.5 mM ethylenediamine-tetraacetic acid, for example). After grinding of the tissue, an equal volume of a 50% (v/v) slurry of washed polyvinylpyrrolidone (suspended in grinding buffer) is added and mixed thoroughly. After centrifugation of the mixture, a sample of the cleared extract is subjected to electrophoresis through a non-denaturing (8%, w/v) polyacrylamide gel. The resulting gel is used to prepare a sandwich with a thin film (<2 mm) of agarose (0.7 % agarose, 0.5 % Sigma medium viscosity carboxymethylcellulose) bonded to "GELBOND" film (FMC Corporation). After incubation for 1.5 hours at 50° C, the agarose film is stained with "CONGO RED" dye

for 30 minutes followed by a 1M NaCl wash. After several minutes, it is possible to visualize cellulase activity as a clear zone within a background of red staining.

Cellulase activity is most commonly assayed in aqueous solution, using a cellulosic substrate and monitoring the reaction for either the release of a chromophore/fluorophore or release of cellobiose ("reducing sugar"). For example, *T. fusca* E2 activity can be measured by incubating a sample of the enzyme in a 0.4 ml reaction containing 1% (w/v) low viscosity carboxymethylcellulose (Sigma C-5678) and 50 mM NaOAc pH 5.5 at 55°C for 2-20 hours. 1.0 ml of DNS solution, see *Irwin et al.* (1993), is then added and the mixture is boiled for 15 minutes. Measurement of absorbance values at 600 nm for each reaction can then be correlated to values determined for a known series of glucose standards to determine the extent of carboxymethylcellulose hydrolysis. For plant extracts, background values are determined by preparing parallel reaction samples which contain no substrate and subtracting this value from that obtained in the presence of 1% carboxymethylcellulose.

For a more complete discussion of cellulase assays, see *Adney et al.* (1994), *Baker et al.* (1992), *Tucker et al.* (1989) and *Irwin et al.* (1993).

Isolation of Cellulase Activity from Plants:

It is most preferred that, where applicable, the enzyme not be purified from the plant material, but rather that the plant material containing the cellulase activity be used directly. This is demonstrated in the Examples, below, where transgenic alfalfa which expressed cellulase activity is added directly to silage materials to further the extent of fermentation.

If isolation of the cellulase activity is desired, this can be accomplished by any means known to the art. For example, the preferred *T. fusca* E2, E3, and CBH I enzymes are taken from thermo-tolerant bacteria. The activity of these enzymes remains unchanged by treatments up to about 55-60°C. Therefore, these enzymes can be isolated by gently heating the plant material in aqueous buffered solution (100 mM Tris/HCl pH 9.0, for example) to precipitate the bulk of plant proteins. The soluble cellulase enzymes

are then recovered and further purified by any means known to the art, including HPLC, affinity chromatography, and the like. To facilitate downstream processing of the enzyme, a purification tag may optionally be incorporated into the expressed cellulase.

Since the above-mentioned enzymes are well-characterized, the preferred purification scheme is based on established protocols already in existence. For example, *T. fusca* E2 from a heat-treated plant extract is further purified by adsorption to a phenyl "SEPHAROSE" column in the presence of 0.8 M ammonium sulfate. Successive column washes using ammonium sulfate concentrations of 0.6 M and 0.3 M in a buffer containing 5 mM KPi, pH 6.0 and 5 mM NaCl are followed by a final wash with 0.1 M KPi, pH 6.0. Elution of E2 is accomplished using 5 mM KPi, pH 6.0. Peak fractions are loaded on a hydroxylapatite column (equilibrated with 1 mM KPi, pH 6.0) and the flow-through fractions collected and pooled. The pooled fractions are loaded on a "Q-SEPHAROSE" column (pre-equilibrated with 10 mM BisTris, pH 6.0) and eluted with a continuous buffer gradient from 5 mM BisTris, pH 6.0 to 150 mM NaCl, 5 mM BisTris, pH 6.0. Peak fractions from the "Q-SEPHAROSE" column are then pooled, concentrated and stored frozen in convenient aliquots. Similarly detailed protocols exist for both *T. fusca* E3 and *T. reesei* CBH I, see *Irwin et al.* (1993).

EXAMPLES

The following Examples are included solely to aid in a more complete understanding of the manufacture and use of the transgenic plants disclosed and claimed herein. The Examples do not limit the scope of the invention in any fashion.

Example 1: Production of Transgenic Alfalfa and Tobacco Which Express Cellulase "E2" of *T. fusca*

Transgenic alfalfa and tobacco plants were produced using the same protocol. Binary vectors carrying recombinant cellulase expression cassettes were transformed into *Agrobacterium tumefaciens* strain LBA 4404, facilitating *Agrobacterium*-mediated

transformation of plant tissue. The construct used is shown in Fig. 1. The gene encoding the E2 cellulase of *Thermomonospora fusca* was obtained as described by Ghangas & Wilson (1988). The E2 gene was modified by PCR using the XbaE2 primer, 5'-GCTCTAGATGAATGATTCTCCGTTC-3' (SEQ. ID. NO: 4) and the "-20 sequencing primer," 5'-TGACCGGCAGCAAAATG-3' (SEQ. ID. NO: 5), (product #1211, New England Biolabs, Inc., Beverly, Massachusetts), resulting in a recombinant gene in which an *Xba I* site (***bold italics***) was incorporated immediately 5' to an introduced start codon (underlined). This start codon precedes the first codon encoding the mature form of the E2 protein (AAT, nt's 348-350 in SEQ. ID. NO: 1). The net effect of these changes is the removal of the bacterial secretion signal peptide (resulting in cytosolic accumulation), the addition of a novel cloning site to facilitate expression cassette construction and the addition of a methionine residue to the N-terminus of the protein compared to the processed mature form of E2 obtained from *T. fusca*.

The cloned E2 gene required no modification at the 3' end as a convenient *Eco RI* restriction site occurs approximately 45 nucleotides 3' to the stop codon.

The preferred expression cassette includes the hybrid "MAC" promoter and the mannopine synthetase terminator. The MAC promoter contains distal elements, including the transcriptional enhancer, of the CaMV 35S promoter (-940 to -90, relative to the mRNA start site), as well as proximal promoter elements derived from the *Agrobacterium* mannopine synthetase promoter (-301 to +65 relative to the mRNA start site). MAC has been reported to result in higher levels of expression than either of the natural promoters (*Comai et al.* (1990).) The expression cassette was cloned into the pCGN1578 binary vector and mobilized into *Agrobacterium*.

Initial tobacco transformants were screened by western blot to determine the level of expression. Levels of expression ranged up to 0.1 to 0.2 % of extracted protein. The mature plants were allowed to self and set seed. One of the initial transformants, designated CT30, was tested further to verify the sexual transmission of the transgene. S1 seeds from this plant were germinated and tested for kanamycin resistance. Leaf samples from kan^R seedlings as well as a W38 control were prepared for western blot

analysis as described previously. The results are depicted in Fig. 3. Each lane contained extract corresponding to 5 mg fresh weight of leaf tissue. In addition, 1 ng of purified E2 enzyme was loaded as a control. Levels of expression were similar to that observed in the parental transformant, demonstrating the stable sexual transfer of this trait. Similar genetic stability was also observed in alfalfa plants transformed with this transgene.

The thermal stability and enzymatic activity of recombinant *T. fusca* E2 was demonstrated using transgenic alfalfa. Samples were prepared for zymogram analysis as described above. As shown in Fig. 5, aliquots of alfalfa extract were treated for 5, 10 and 20 minutes (grouped from left to right) at each of the temperatures indicated (°C) before being subjected to native gel electrophoresis. An untreated sample of extract and two purified E2 standards were included as controls. Levels of E2 activity corresponded well with expected activity based on western blot analysis of samples from the same plant. In addition, no significant loss in band intensity (activity) was observed at any of the treatment temperatures, despite the fact that greater than 95% of the soluble protein in the extract is denatured after 20 minutes at 65°C.

Example 2: Production of Transgenic Alfalfa and Tobacco Which Express Cellulase "E3" of *T. fusca*

Here, the expression construct depicted in Fig. 2 was used to transform alfalfa and tobacco using the same methodology as described in Example 1.

The gene encoding the E3 cellulase of *Thermomonospora fusca* was obtained as described by Zhang *et al.* (1995). The 5' end of the E3 gene was modified by PCR using the primer XbaE3, 5'-GCTCTAGATGGCCGGCTGCTCGGTG-3' (SEQ. ID. NO: 6), resulting in a recombinant gene in which an *Xba I* site (***bold italics***) was incorporated immediately 5' to an introduced start codon (underlined). This start codon precedes the first codon encoding the mature form of the E3 protein (GCC, nt 689-691 in SEQ. ID. NO: 2). The 3' end of the E3 gene was modified using the primer RIE3, 5'-GGAATTCTTACAGAGGCGGGTAG-3' (SEQ. ID. NO: 7), thereby placing an *Eco RI*

restriction site (***bold italics***) immediately 3' to the stop codon (underlined) for the E3 gene. Note that this latter primer is homologous to the noncoding strand of the E3 gene. The net effect of these changes is the removal of the bacterial secretion signal peptide (resulting in cytosolic accumulation), the addition of novel cloning sites to facilitate expression cassette construction and the addition of a methionine residue to the N-terminus of the protein (compared to the processed, mature form of E3 obtained from *T. fusca*).

The E3 expression cassette was constructed as described above for the E2 cassette.

Initial tobacco transformants were screened by western blot to determine the level of expression. Levels of expression ranged up to about 0.04% of extracted protein. The mature plants were allowed to self and set seed. One of the initial transformants, designated CT117, was tested further to verify the sexual transmission of the transgene. SI seeds from this plant were germinated and tested for kanamycin resistance. Leaf samples from kan^R seedlings as well as a W38 control were prepared for western blot analysis as described previously. The results are depicted in Fig. 4. Each lane contained extract corresponding to 5 mg fresh weight of leaf tissue. In addition, 1 ng of purified E3 enzyme was loaded as a control. Levels of expression were comparable to that observed in the parental transformant, demonstrating the stable sexual transfer of this trait.

Example 3: Sexual Transfer of Cellulase Expression in Tobacco and Alfalfa

Original transgenic lines of tobacco and alfalfa shown to express either E2 or E3 cellulase were used in sexual crosses. In both cases, the trait segregated in progeny as predicted by Mendelian genetics. Expression levels were the same as, or greater than those seen in parental lines.

Example 4: Use of Transgenic Alfalfa in Silage

Here, regular non-transformed alfalfa, alfalfa transformed according to Example 1 and alfalfa transformed according to Example 2 were ensiled under identical conditions

for one month and the products of fermentation for each experiment quantified. The results are presented in Table 2.

All of the ensiled plant material was ground separately through a manual meat grinder. The grinder was rinsed with water and wiped with ethanol after grinding each sample. A 1 to 1 to 1 mixture of non-transformed alfalfa, E2-transformed alfalfa, and E3-transformed alfalfa was ground together and used to assemble two control silos (Cont1 and Cont2, 50 g each). The two control silos were inoculated with a 1 mL of a commercial inoculant (0.1098 g "BIOMATE LP/PC" concentrate in 500 mL sterile water).

Two silos each of E2-transformed alfalfa (E2-1, E2-2) and E3-transformed alfalfa (E3-1, E3-2) were constructed in the same fashion as the controls (35 g each, inoculated with 0.6 mL of the above-noted inoculant). Two silos of mixed E2- and E3-transformed alfalfa were constructed by grinding together 17.5 g each of E2- and E3-transformed alfalfa per silo (35 g each, inoculated with 0.6 mL of the above-noted inoculant).

All of the silos were then placed into a 30°C water bath until opening.

Of special note in this Example is the increased amount of fermentation products in the transgenic alfalfa as compared to the non-transformed alfalfa. In particular, note that a mixture of alfalfa herbage expressing both the E2 and E3 cellulases exhibits markedly improved fermentation yield as compared to the non-transformed alfalfa and ensiled alfalfa expressing either E2 or E3 enzymes.

Clearly, as shown by this Example, expression of cellulases in transgenic alfalfa leads to better silage production.

TABLE 2

							Organic Acid Analysis (OAA) via HPLC							
Sample														
ID	%DM	pH	SUC	LAC	FOR	ACE	PRO	2,3But	ETOH	BUT	Total Prod.	Avg. Total Prod.		
Cont 1	23.33612	5.741	0.144	2.014	0.000	1.764	0.000	0.239	0.704	0.000	4.86	5.47		
Cont 2	21.62983	5.121	0.159	3.885	0.000	1.407	0.328	0.000	0.294	0.000	6.07			
E2-1	22.99369	5.277	0.157	3.390	0.000	2.300	0.233	0.000	0.330	0.000	6.41	6.09		
E2-2	23.83774	5.166	0.361	2.935	0.000	1.998	0.177	0.000	0.298	0.000	5.77			
E3-1	22.88773	5.128	0.283	3.321	0.000	2.380	0.177	0.000	0.292	0.000	6.45	6.61		
E3-2	22.22822	5.151	0.354	3.324	0.000	2.608	0.200	0.000	0.288	0.000	6.77			
E23-1	22.95945	5.743	0.551	2.848	0.000	3.185	0.328	0.000	0.337	0.000	7.25	7.45		
E23-2	22.66411	5.888	0.602	2.745	0.000	3.649	0.301	0.000	0.353	0.000	7.65			

The table headings are as follows:

%DM = percent dry matter of silage, pH = acidity, SUC = succinic acid, LAC = lactic acid, FOR = formic acid, ACE = acetic acid, PRO = propionic acid, 2,3 But = 2,3-butanediol, ETOH = ethanol, BUT = butyric acid, Cont 1 and Cont 2 = controls, E2-1 and E2-2 = transformant expressing E2, E3-1 and E3-2 = transformants expressing E3, E23-1 and E23-2 = a 1:1 mixture of herbage from transgenic alfalfa expressing E2 and E3.

Example 5: Production of Transgenic Tobacco Plants Which Express Cellulase "E1" of *A. cellulolyticus*

Transgenic tobacco plants were produced in a manner analogous to that described in Examples 1 and 2. Binary vectors carrying recombinant cellulase expression cassettes were transformed into *Agrobacterium tumefaciens* strain LBA 4404, thereby facilitating *Agrobacterium*-mediated transformation of plant tissue. The constructs for this Example are depicted schematically in Figs. 6A and 6B. The gene encoding the E1 cellulase of *Acidothermus cellulolyticus* was obtained as described previously by *Himmel et al.* (see U.S. Patent No. 5,275,944). The E1 gene was then modified by PCR using the *NarE1* primer, 5'-CGGGGCGCCGGCGGGCGGCTAT-3' (SEQ. ID. NO: 10) and the *SacE1* primer, 5'-CCGAGCTCTTAAGTTGCTGC-3' (SEQ. ID. NO: 11) to generate a recombinant E1 gene. The recombinant gene has a *NarI* site at the 5' end and a *SacI* site at the 3' end (restriction sites are underlined) to facilitate fusion to the VSP leader coding sequence (SEQ. ID. NO: 3) and nopaline synthetase terminator.

As in the previous Examples, PCR-derived fragments were sequenced to verify that no errors (mutations) had been introduced. The resulting cassette includes the VSP leader sequence operationally linked to the "mature" portion of the E1 coding sequence. This cassette was then cloned into pBI121 (Clontech Labs, Palo Alto, California) as a *XbaI* to *SacI* fragment, replacing the *uidA* gene and placing the new construct (designated pZ49.1) under the control of the CaMV 35S promoter.

An analogous construct (designated pZ57.1) was generated in which the E1 coding sequence was truncated to yield the E1 catalytic domain (E1cd) using the *NarE1* primer (SEQ. ID. NO: 10) and the *SacE1cd* primer, 5'-TGGAGCTCTAGACAGGATCGAAAAT-3' (SEQ. ID. NO: 12). This construct encodes a polypeptide containing the VSP leader peptide (SEQ. ID. NO: 13) fused to the first 358 amino acids of the E1 protein. The codon specifying valine 358 is bold, italics (note that this oligonucleotide represents the "antisense" strand). Plasmids pZ49.1 and pZ57.1 were transformed into *Agrobacterium tumefaciens* strain LBA4404 to yield strains PZA8 and PZA9, respectively.

Putative transgenic tobacco plants were screened by a combination of Western blotting and E1 activity assay. Leaf samples were removed from plants grown in Magenta boxes (MS medium). Samples were ground in E1 grinding buffer (50 mM NaOAc pH 5.5, 100 mM NaCl, 10% (v/v) glycerol, 1 mM ethylenediamine-tetraacetic acid, 1 mM phenylmethanesulfonyl fluoride, 1 mg/l aprotinin, 1 mg/l leupeptin, 1 mg/l pepstatin), added at a ratio of 2 μ l per mg of sample. Samples of extract were centrifuged at >10,000 x G for 5 minutes to remove insoluble material and diluted 100-fold in E1 grinding buffer to which acetylated bovine serum albumin had been added (0.1 mg/ml final concentration). Diluted extract was assayed for activity at 65°C using β -D-cellobiopyranoside (MUCB) as a substrate (0.5 mM MUCB, 50 mM NaOAc pH 5.5, 100 mM NaCl).

Reactions were terminated by the addition of an equal volume of 150 mM glycine/NaOH (pH 10). Fluorescence at 460 nm was quantified using a commercial plate reader ("BIOLUMIN 960," Molecular Dynamics) with excitation set at 355 nm. Enzyme activity in extracts was then compared to the activity of purified E1 holoenzyme and E1 catalytic domain (generously provided by Steve Thomas, National Renewable Energy Laboratory, Golden, Colorado).

In addition, a set of 4-methylumbelliferone standards was also assembled for use as calibration standards. The same extracts were also subjected to analysis by Western blotting. Both PZA8- and PZA9-transformed tobacco plants accumulated an immunoreactive species that co-migrates with purified E1cd. Very little full-length E1 is present in PZA8 transformants, indicating that proteolytic processing of the E1 enzyme is taking place. For this reason, activities are reported as E1cd equivalent, even in those plants that contain an intact E1 coding sequence. For PZA8 transformants, the average E1 expression level was 0.10% of total soluble protein, with the highest expressing plant accumulating E1 at 0.33% of total soluble protein (see Fig. 8A). E1 expression was higher in PZA9 transformants, with an average expression level of 0.21% of total soluble protein and a high value of 0.59% (See Fig. 8B).

Example 6: Production of Transgenic Tobacco Plants Which Express Cellobiohydrolase I (CBH I) of *Trichoderma reesei*

Transgenic plants were generated essentially as described in Example 5. A schematic diagram of the construct used is shown in Fig. 7. The gene encoding CBH I of *Trichoderma reesei* (SEQ. ID. NO: 9) was generously provided by Steve Thomas (National Renewable Energy Laboratory, Golden, Colorado) and is substantially the same as the gene described by *Shoemaker et al.* with its introns removed. Sequence data obtained by the inventors and by NREL scientists indicates that the gene used in this Example differs from the *Shoemaker et al.* sequence at nucleotide 1429. Specifically, the gene used here contains a 4 base-pair sequence (CGCC) inserted in place of G1429, thereby effectively inserting an additional codon and replacing Arg459 with two prolines. A similar substitution exists in a related CBH I enzyme from *Trichoderma viride* (see *Cheng et al.*, 1990). The gene used here also has a silent mutation in the codon specifying Thr41 (ACT changed to ACG).

The CBH I gene was modified by PCR using the cbh2-2 primer, 5'-GCTCTAGATGTATCGGAAGTTGGC-3' (SEQ. ID. NO: 14) and the cbh3-1 primer, 5'-CCCCCGGGTTACAGGCACTGAGAG-3' (SEQ. ID. NO: 15) to generate a recombinant CBH I gene which retains its secretory leader peptide. The recombinant gene has an *Xba*I site at the 5' end and a *Xma*I site at the 3' end (restriction sites are shown in underline, start codon is bold, italics) to facilitate vector construction. The gene was cloned into pBI121 (Clontech) as a *Xba*I to *Xma*I fragment, replacing the *uidA* gene and placing the CBH I gene under the control of the CaMV 35S promoter.

Putative transgenic tobacco plants were screened by Western blotting. Leaf samples were removed from plants grown in Magenta boxes (MS medium) and ground in E1 grinding buffer (see composition in Example 5), added at a ratio of 2 μ l per mg of sample. The extract was centrifuged at >10,000 x G for 5 minutes to remove insoluble material and a portion prepared for SDS-PAGE and subsequent blotting and detection. See Fig. 9, which depicts the SDS-PAGE gel. Of 12 plants screened in this way, 3 had detectable expression (about 0.01% of total soluble protein). In addition,

immunoreactive material migrated slightly ahead of the RuBisCo large subunit, consistent with the expected mobility of the 48 kDa catalytic domain (Divne *et al.*, 1994). This suggests that CBH I, like *A. cellulolyticus* E1, is proteolytically cleaved by a plant protease.

5

Example 7: Production of Transgenic Tobacco Plants Which Express the Endoglucanase Encoded by the *cenA* Gene of *Cellulomonas fimi*

In a manner analogous to the previous Examples, the *cenA* gene of *Cellulomonas fimi* (Wong *et al.*, 1986) can be expressed in tobacco. A schematic diagram of the construct used is shown in Fig. 10. The *cenA* gene (SEQ. ID. NO: 16) is modified by PCR using the *cenA*pst primer, 5'-GGCTGCAGGCGGCTGCCGCGTCGAC-3' (SEQ. ID. NO: 17) and the *cenA*sac primer, 5'-CCGAGCTCTCACCACCTGGCGTT-3' (SEQ. ID. NO: 18) to generate a recombinant *cenA* gene. The recombinant gene has a *Pst*I site at the 5' end and a *Sac*I site at the 3' end (restriction sites are underlined, novel glycine codon in bold italics) to facilitate fusion to a VSP leader coding sequence (SEQ. ID. NO: 19) and nopaline synthetase terminator. In addition, the proline at position 2 in the mature endoglucanase enzyme is changed to a glycine, a conservative substitution. The resulting cassette consists of the VSP leader sequence fused to the "mature" portion of the *cenA* gene.

This cassette is then cloned into pBI121 (Clontech) as a *Xba*I to *Sac*I fragment, replacing the *uidA* gene and placing the new construct under the control of the CaMV 35S promoter. As in the previous Examples, *Agrobacterium tumefaciens* strain LBA4404 is transformed with the resulting binary vector and subsequently used to transform plants.

Example 8: Production of Transgenic Tobacco Plants Which Express Endoglucanase D, Encoded by the *celD* Gene of *Clostridium thermocellum*

In a manner analogous to the previous Examples, the *celD* gene of *Clostridium thermocellum* (see Joliff *et al.*, 1986) can be expressed in tobacco. A schematic diagram

of the construct used is shown in Fig. 11. The *celD* gene (SEQ. ID. NO: 20) is modified by PCR using the *celD*_{pst} primer, 5'-AGCTGCAGAAATAACGG-3' (SEQ. ID. NO: 21) and the *celD*_{sac} primer, 5'-CCGAGCTCTTATATTGGTAATTTCTC-3' (SEQ. ID. NO: 22) to generate a recombinant *celD* gene. The recombinant gene has a *Pst*I site at the 5' end and a *Sac*I site at the 3' end (restriction sites are underlined) to facilitate fusion to the VSP leader coding sequence (SEQ. ID. NO: 19) and nopaline synthetase terminator. The resulting cassette includes the VSP leader sequence fused to the "mature" portion of the *celD* gene. Subsequent manipulations are be carried out as described in the previous Examples.

Example 9: Production of Transgenic Tobacco Plants Which Express Exoglucanase S, Encoded by the *exgS* Gene of *Clostridium cellulovorans*

In a manner analogous to the previous Examples, the *exgS* gene of *Clostridium cellulovorans* (see Liu and Doi, 1998) can be expressed in tobacco. A schematic diagram of the construct used is shown in Fig. 12. The *exgS* gene (SEQ. ID. NO: 23) is modified by PCR using the *exgS*_{nar} primer, 5'-CGGGGCGCCGACACAGTAGTGCCA-3' (SEQ. ID. NO: 24) and the *exgS*_{sac} primer, 5'-CCGAGCTCTTATTTAATCTTAAGC-3' (SEQ. ID. NO: 25) to generate a recombinant *exgS* gene. The recombinant gene has a *Nar*I site at the 5' end and a *Sac*I site at the 3' end (restriction sites are underlined) to facilitate fusion to the VSP leader coding sequence (SEQ. ID. NO: 13) and nopaline synthetase terminator. The resulting cassette consists of the VSP leader sequence fused to the "mature" portion of the *exgS* gene. Subsequent manipulations are carried out as described previously.

Example 10: Production of Transgenic Tobacco Plants Which Express Exocellulase E6, Encoded by the *celF* Gene of *Thermobifida fusca* (formerly *Thermomonospora fusca*)

In a manner analogous to the previous Examples, the *celF* gene of *Thermobifida fusca* (see Irwin et al., 1999) can be expressed in tobacco. A schematic diagram of the

construct used is shown in Fig. 13. The *celF* gene (SEQ. ID. NO: 26) is modified by PCR using the *celF*pst primer, 5'-ACGCTGCAGTCGCCTGCTCGG-3' (SEQ. ID. NO: 27) and the *celF*xma primer, 5'-CCCCCGGGTCAGGGAGCTCCGGC-3' (SEQ. ID. NO: 28) to generate a recombinant *celF* gene. The recombinant gene has a *Pst*I site at the 5' end and a *Xma*I site at the 3' end (restriction sites are underlined) to facilitate fusion to the VSP leader coding sequence (SEQ. ID. NO: 19) and nopaline synthetase terminator.

The *celF* gene itself contains two internal *Xma*I recognition sites, which are removed by site-directed mutagenesis. Briefly, a portion of the gene containing the sites is subcloned to pBluescript KS+ (Stratagene, La Jolla, California) as a *Bgl* II to *Xho* I fragment. PCR reactions are carried out using primer 2777 (5'-GGCCACCTGGGCAGG-3', SEQ. ID. NO: 29) and the M13-20 sequencing primer (5'-GTAAAACGACGGCCAGT-3', SEQ. ID. NO: 30), thereby destroying the site at 2775 in the Genbank sequence (underline indicates mutated nucleotide).

Similarly, primer 3227 (5'-GCGACGCTCGGGCCG-3', SEQ. ID. NO: 31) and the reverse sequencing primer (5'-AACAGCTATGACCATG-3', SEQ. ID. NO: 32) destroy the site at 3227. The two overlapping amplified fragments are then purified, heated briefly to 95°C and cooled gradually to allow annealing to occur. The annealed template is subjected to another round of PCR using the M13-20 sequencing primer (SEQ. ID. NO: 30) and the reverse sequencing primer (SEQ. ID. NO: 32). This fragment is then subcloned as a *Bgl* II to *Xho* I fragment and sequenced before being used to replace the wild-type *celF* sequence. Both base changes are at the 3rd position in the codon and do not alter protein sequence. The resulting cassette consists of the VSP leader sequence operationally linked to the "mature" portion of the *celF* gene. Subsequent manipulations are carried out as described hereinabove.

The invention is not limited to the preferred embodiments, transformation protocols, transformed plant hosts, and expression constructs explicitly described above, but encompasses all such forms thereof as are encompassed within the scope of the attached claims.

BIBLIOGRAPHY

- Adney et al.* (1994), Cellulase assays. In: *Enzymatic conversion of biomass for fuels production*, Eds. M. E. Himmel, J.O. Baker & R.P. Overend. ACS symposium series 566.
- Aspegren et al.* (1995), Secretion of a heat-stable fungal β -glucanase from transgenic, suspension-cultured barley cells. *Molecular Breeding* 1:91-99.
- Baker et al.* (1992), Thermal denaturation of *T. reesei* cellulases studied by differential scanning calorimetry and tryptophan fluorescence. *Apply. Biochem. Biophys.* 34:217-231.
- Bednarek* (1991), The barley lectin carboxy-terminal peptide is a vacuolar protein sorting determinant in plants. *The Plant Cell* 3:1195-1206.
- Belkacemi et al.* (1996), Enzymatic hydrolysis of timothy grass pretreated by ammonia fiber explosion. In: *Liquid fuels and industrial products from renewable resources, Proceedings of the third liquid fuel conference*, Eds. J.S. Cundiff, E.E. Gavett, C. Hansen, C. Peterson, M.A. Sanderson, H. Shapouri & D.L. VanDyne. ASAE publication 08-96 pp 232-240.
- Bingham et al.* (1975), Breeding alfalfa which regenerates from callus tissue in culture. *Crop Sci.* 15:719-721.
- Brown and Atanassov* (1985), Role of genetic background in somatic embryogenesis in *Medicago*. *Plant Cell Tissue Organ Culture* 4:107-114.
- Carrer et al.* (1993), Kanamycin resistance as a selectable marker for plastid transformation in tobacco. *Mol. Gen. Genet.* 241:49-56.
- Castillo et al.* (1994), Rapid production of fertile transgenic plants of Rye. *Bio/Technology* 12:1366-1371.
- Cheng et al.* (1990), *Nucl. Acids Res.* 18:5559.
- Comai et al.* (1990), Novel and useful properties of a chimeric plant promoter combining CaMV 35S and MAS elements. *Plant Mol. Biol.* 15:373-381.
- Coughlan, M.P.* (1988), Staining Techniques for the Detection of the Individual Components of Cellulolytic Enzyme Systems. *Methods in Enzymology* 160:135-144.

Current Protocols in Molecular Biology, Volumes 1-3, Series Editor, Virginia Benson Chanda, ©1987-1997, John Wiley & Sons, Inc.

de Castro Silva Filho et al. (1996), Mitochondrial and chloroplast targeting sequences in tandem modify protein import specificity in plant organelles. *Plant Mol. Biol.* 30:769-780.

Divne et al. (1994), The three-dimensional crystal structure of the catalytic core of cellobiohydrolase I from *Trichoderma reesei*. *Science* 265:524-528.

Ghangas & Wilson (1988), Cloning of the *Thermomonospora fusca* endoglucanase E2 gene in *Streptomyces lividans*: Affinity purification and functional domains of the cloned gene product. *Appl. Envir. Microbiol.* 54:2521-2526.

Grohmann et al. (1992), Potential for fuels from biomass and wastes. In: *Emerging technologies for materials and chemicals from biomass*, Eds. R.M. Powell, T.P. Schultz and R. Narayan. ACS symposium series 576.

Henrissat et al. (1995), Synergism of cellulases from *Trichoderma reesei* in the degradation of cellulose. *Bio/Technology* 3:722-726.

Horsh et al. (1985), A simple and general method for transferring genes into plants. *Science* 227:1229-1231.

Irwin et al. (1993), Activity studies of eight purified cellulases: Specificity, synergism, and binding domain effects. *Biotechnol. Bioeng.* 42:1002-1013.

Irwin et al. (1999), Characterization of a *Thermomonospora fusca* family 48 exocellulase E6. Direct Genbank submission AF144563.

Ishida et al. (1996), High efficiency transformation of maize mediated by *Agrobacterium tumefaciens*. *Nature Biotechnology* 14:745-750.

Joliff et al. (1986), Nucleotide sequence of the cellulase gene *celD* encoding endoglucanase D of *Clostridium thermocellum*. *Nucleic Acids Res.* 14:8605-8613.

Keegstra et al. (1993), Targeting of proteins into chloroplasts. *Physiologia Plantarum* 93:157-162.

Lao et al. (1991), *J. Bacteriol.* 173:3397-3407.

Liu and Doi (1998), Properties of *exgS*, a gene for a major subunit of the *Clostridium cellolovorans* cellulosome. *Gene* 211:39-47.

- Mason et al.* (1988), Proteins homologous to leaf glycoproteins are abundant in stems of dark-grown soy bean seedlings. Analysis of proteins and cDNAs. *Plant Mol. Biol.* 11:845-856.
- McBride and Summerfelt* (1990), Improved binary vectors for *Agrobacterium* mediated plant transformation. *Plant Mol. Biol.* 14:269-276.
- McBride et al.* (1994), Controlled expression of plastid transgenes in plants based on a nuclear DNA-encoded and plastid-targeted T7 RNA polymerase. *Proc. Natl. Acad. Sci. USA* 91:7301-7305.
- Micelli et al.* (1996), Integrated treatments of steam explosion and enzymatic hydrolysis to produce energetic and industrial products from lignocellulosic biomasses. *Agro-food-Industry Hi-tech* 7:25-28.
- Murashige and Skoog* (1962), A revised medium for rapid growth and bioassays with tobacco tissue cultures. *Physiol. Plant* 15:473-497.
- Pentilla et al.* (1987), *Yeast* 3:175-185.
- Shoemaker et al.* (1983), *Bio/Technology* 1:691-696.
- Sonnewald et al.* (1991), Transgenic tobacco plants expressing yeast-derived invertase in either the cytosol, vacuole or apoplast: a powerful tool for studying sucrose metabolism and sink/source interactions. *The Plant J.* 1:95-106.
- Spezio et al.* (1993), Crystal structure of the catalytic domain of a thermophilic endocellulase. *Biochemistry* 32:9906-9916.
- Tucker et al.* (1989), Ultra-thermostable cellulases from *Acidothermus cellulolyticus* comparison of temperature optima with previously reported cellulases. *Biotechnology* 7:817-820.
- Vasil et al.* (1993), Rapid production of transgenic wheat plants by direct particle bombardment of cultured immature embryos. *Bio/Technology* 11:1553-1558.
- Wandelt et al.* (1992), Vicilin with carboxy-terminal KDEL is retained in the endoplasmic reticulum and accumulates to high levels in the leaves of transgenic plants. *Plant J.* 2:181-192.
- Wong et al.* (1986), Characterization of an endoglucanase gene *cenA* of *Cellulomonas fimi*. *Gene* 44:315-324.

Zambryski, P., J. Tempe, and J. Schell (1989), Transfer and function of T-DNA genes from *Agrobacterium* Ti and Ri plasmids in plants. *Cell* 56:193-201.

Zhang *et al.* (1995), Characterization of a *Thermomonospora fusca* exocellulase. *Biochemistry* 34:3386-3395.

CLAIMS

What is claimed is:

1. A genetically recombinant plant comprising a plant transformed to contain and express in recoverable quantities an exogenous gene sequence or fragment thereof which encodes a cellulose-degrading gene product and wherein the gene product is classified within an enzyme classification selected from the group consisting of EC 3.2.1.4, EC 3.2.1.6, EC 3.2.1.21, EC 3.2.1.91, and combinations thereof.
2. The genetically recombinant plant of Claim 1, wherein the gene product is expressed constitutively.
3. The genetically recombinant plant of Claim 1, wherein the gene product is expressed stage-specifically.
4. The genetically recombinant plant of Claim 1, wherein the gene product is expressed tissue-specifically.
5. The genetically recombinant plant of Claim 4, wherein the gene product is expressed in a plant tissue selected from the group consisting of seeds, fruit, stems, leaves, and tubers.
6. The genetically recombinant plant of Claim 4, wherein the plant contains at least two exogenous genes and wherein their respective gene products are expressed independently of one another.
7. The genetically recombinant plant of Claim 1, wherein the gene product is expressed in a targeted sub-cellular compartment.

8. The genetically recombinant plant of Claim 7, wherein the gene product is expressed in a sub-cellular compartment selected from the group consisting of: plastid, cytosol, endoplasmic reticulum, mitochondrion, inclusion body, and vacuole.
9. The genetically recombinant plant of Claim 7, wherein the plant contains at least two exogenous genes and wherein their respective gene products are expressed independently of one another.
10. The genetically recombinant plant of Claim 1, wherein the gene product is expressed extra-cellularly.
11. The genetically recombinant plant of Claim 1, wherein the plant is dicotyledonous.
12. The genetically recombinant plant of Claim 1, wherein the plant is monocotyledonous.
13. The genetically recombinant plant of Claim 1, wherein the plant expresses a cellulose-degrading gene product classified within an enzyme classification selected from the group consisting of EC 3.2.1.4 and EC 3.2.1.91.
14. The genetically recombinant plant of Claim 13, wherein the plant expresses *A. cellulolyticus* endoglucanase E1 or *T. reesei*. CBH I.
15. The genetically recombinant plant of Claim 1, which is alfalfa or tobacco.
16. The genetically recombinant plant of Claim 15, which is stably transformed to contain a gene sequence which encodes a cellulase-degrading enzyme selected from the group consisting of *T. fusca* cellulase E2, *T. fusca* cellulase E3, *T.*

ressei CBH I, *A. cellulolyticus* endoglucanase E1, and combinations thereof.

17. The genetically recombinant plant of Claim 16, which is alfalfa.
18. The genetically recombinant plant of Claim 16, which is tobacco.
19. The genetically recombinant plant of Claim 16, which is alfalfa transformed to contain and express a gene sequence selected from the group consisting of SEQ. ID. NOS: 8 and 9.
20. The genetically recombinant plant of Claim 16, which is tobacco transformed to contain and express a gene sequence selected from the group consisting of SEQ. ID. NOS: 8 and 9.
21. A method for producing cellulose-degrading enzymes comprising cultivating a genetically recombinant plant according to Claim 1.
22. The method of Claim 21, further comprising concentrating the cellulose-degrading enzymes.
23. A method for producing cellulose-degrading enzymes comprising cultivating a genetically recombinant plant according to Claim 13.
24. The method of Claim 23, further comprising concentrating the cellulose-degrading enzymes.
25. A method of ensiling comprising ensiling a plant according to Claim 1, whereby cellulose-degrading enzymes produced by the plant increase nutritional value of silage.

26. A method of ensilement comprising ensiling a plant according to Claim 13, whereby cellulose-degrading enzymes produced by the plant increase nutritional value of silage.

TRANSGENIC PLANTS AS AN ALTERNATIVE SOURCE OF LIGNOCELLULOSIC-DEGRADING ENZYMES

ABSTRACT OF THE DISCLOSURE

Transgenic plants which express cellulose-degrading enzymes, methods to make the transgenic plants, and methods to use the cellulose-degrading enzymes produced by the transgenic plants are disclosed.

660480.022260

E2 W38 _____ S1 seedlings




FIG. 3

E3 W38 — S1 seedlings




FIG. 4

666180" 666180" 666180"




FIG. 5




FIG. 6a




FIG. 6b






FIG. 7






FIG. 9




FIG. 10




FIG. 11




FIG. 12




FIG. 13

SEQUENCE LISTING

<110> Austin-Phillips, Sandra
Burgess, Richard D
German, Thomas L
Ziegelhoffer, Thomas

<120> Transgenic Plants as an Alternative Source of
Lignocellulosic-Degrading Enzymes

<130> Transgenic Plants Expressing Cellulase

<140>

<141>

<150> 08/883,495

<151> 1997-06-26

<160> 32

<170> PatentIn Ver. 2.0

<210> 1

<211> 1621

<212> DNA

<213> Thermomonospora fusca

<400> 1

```
cgatatggat gatctgacgt ctgaatcccc ttgtcaccct agacattcac ccattttgtc 60
gctttttacgg cttttcttgg gagttctccg tttcaccaag gaacaaaacc gcaacggaga 120
gtaggcgcggt tctttacagc tcccttgcca atggttatcg tccgaacgga aaacgatctg 180
ggagcgctcc cagccatgcg ctctcttcg tgccccctcac ttcttttgag ccttgtgtctc 240
gttaggagcc ccgaatgtcc ccagacctc ttgcgcgtct tctgggcgcc gcggcgggcg 300
ccttggtcag cgcggctgct ctggccttcc ggtcgcaagc ggcgccaat gattctccgt 360
tctacgtcaa cccaacatg tcttcgcgcg aatgggtgcg gaacaacccc aacgacccgc 420
gtaccccggt aatccgcgac cggatcgcca gcgtgcgcga gggcacctgg ttgcgccacc 480
acaaccccgg gcagatcacc ggccagatcg acgcgctcat gagcgccgcc caggccgcgc 540
gcaagatccc gatcctgggt gtgtacaacg ccccgggccg cgactgcggc aaccacagca 600
gcggcgggcg cccagtcac agcgctacc ggtcctggat cgacgaattc gctgcgggac 660
tgaagaaccg tcccgccac atcatcgctg ggccggacct gatctcgctg atgtcgagct 720
gcatccagca cgtccagcag gaagtcctgg agacgatggc gtacgcgggc aaggccctca 780
aggccgggtc ctgcaggcg cggatctact tcgacgccg ccaactccgc tggcactcgc 840
ccgcacagat ggcttctctg ctccagcagg ccgacatctc caacagcgcg cacggtatcg 900
ccaccaaacac ctccaactac cgggtggacc ctgacgaggt cgcctacgcc aaggcggtgc 960
tctcggccat cggcaacccg tccctgcgcg cggtcacga caccagccgc aacggcaacg 1020
gccccgcggg taacgagtgg tgcgaccca gcggacgcgc catcggcacg cccagcacca 1080
ccaacaccgg cgaccgatg atcgacgct tctgtggat caagctgccg ggtgaggccg 1140
acggctgcat cgcgggcgcc ggccagttcg tccgcaggc ggcctacgag atggcgatcg 1200
```

```

ccgcggggcg caccaacccc aaccogaacc ccaacccgac gccaccccc actccgaccc 1260
ccacgccgcc tcccggtcc tggggggcgt gcacggcgac gtacacgac gccaacgagt 1320
ggaacgacgg cttccaggcg accgtgacgg tcaccgcgaa ccagaacatc accggctgga 1380
ccgtgacatg gaccttcacc gacggccaga ccatcaccaa cgcctggaac gccgacgtgt 1440
ccaccagcgg ctccctcggtg accgcgcgga acgtcggcca caacggaacg ctctcccagg 1500
gagccccac agagttcggc ttctgctggct ctaagggcaa ctccaactct gttccgaccc 1560
ttacctgcgc cgccagctga cccctcctgg cagtgcactg ggtggcttag gcgtgctggg 1620
g
1621

```

<210> 2

<211> 3503

<212> DNA

<213> Thermomonospora fusca

<400> 2

```

cggcgatccc cctcatcatt caggtgcggt tagttcccc aggctaccga ggaccgaatt 60
tcggtccggt tttcttgagg cgagccctga gaccgtttcc tgttccgttc cgtcaccatc 120
cttgccgctc ccggcggagg ggggaagcac cccgcgagat ggctccgcca cggcctgttt 180
ccgacccccg tcacaaaagc ccatttaacg cggatattac aaccggtcat gaagtggcta 240
ctctcttttg ggagcgctcc cgtgccgcta gtcacactgg gacgtgaatg gcgtcacggg 300
agggctcgtc gtgtgacaac cattttcgac cctgctttaa gtccctaagt gggagcgctc 360
ccagccttcg ggagaactcc cacacaacca accgtccgac gccactctcc cagcgctcaa 420
acggaggcag cagtgttcac catccccgc tccctccgg ggcccccggc cgtcgctccg 480
gcaaccaccc cgaccggtcg gctgaacact gcagcgctcc gttctcgacc atccccctgc 540
gagagaacat cctccaacca aggaagacac cgatatgagt aaagttcgtg ccacgaacag 600
acgttcgtgg atgcggcgcg ggctggcagc cgctctgga ctggcgcttg gcgcctccat 660
ggtggcgctt gctgctccgg ccaacgcgc cggtgctcg gtgtactaca cgggtcaactc 720
ctggggtaac ggggttaacc ccaacgtcac catcaccaac ctcggcagtg cgatcaacgg 780
ctggaccctg gagtgggact tccccggcaa ccagcaggtg accaacctgt ggaacggggc 840
ctacacccag tccgggcagc acgtgtcggt cagcaacgcc ccgtacaacg cctccatccc 900
ggccaacgga acggttgagt tcgggttcaa cggtcctac tcgggcagca acgacatccc 960
ctcctccttc aagctgaacg gggttacctg cgacggctcg gacgaccccg accccgagcc 1020
cagccccctc ccagccctt cccccagccc cacagacccg gatgagccgg gcggcccagc 1080
caaccgcgcc accaaccocg gcgagaaggt cgacaaccgg ttcgagggcg ccaagctgta 1140
cgtgaaccoc gtctggctcg ccaaggccgc cgtgagccg ggcggttccg cggtcgccaa 1200
cgagtccacc gctgtctggc tggaccgtat cggcgggac gagggcaacg acagcccagc 1260
cacgggctcc atgggtctgc gcgaccacct ggaggaggcc gtccgccagt ccggtggcga 1320
cccgctgacc atccaggctc tcatctacca cctgccggc cgcgactcgc ccgcgtggc 1380
ctccaacggg gagctgggtc ccgatgaact cgaccgctac aagagcgagt acatcgaccc 1440
gatcgccgac atcatgtggg acttcgcaga ctacgagaac ctgcggatcg tcgcatcat 1500
cgagatcgac tccctgccc acctcgtcac caacgtgggc gggaacggcg gcacogagct 1560
ctgcgcctac atgaagcaga acggcggcta cgtcaacggg ttcggctacg cctccgcaa 1620
gctgggcgag atcccgaacg tctacaacta catcgacgcc gccaccacg gctggatcgg 1680
ctgggactcc aacttcggcc cctcgggtga catctctac gaggccgcca acgcctccgg 1740
ctccaccgtg gactacgtgc acggcttcat ctccaacacg gccaaactact cggccactgt 1800
ggagccgtac ctggacgtca acggcacctg taacggccag ctcatccgcc agtccaagtg 1860
ggttgactgg aaccagtagc tcgacgagct ctcttcgtc caggacctgc gtcaggccct 1920
gatcgccaag ggcttcgggt ccgacatcgg tatgtctatc gacacctcc gcaacggctg 1980

```

```

gggtggcccg aaccgtccga ccggaccgag ctccctccacc gacctcaaca cctacgttga 2040
cgagagccgt atcgaccgcc gtatccaccc cggtaactgg tgcaaccagg ccggtgcggg 2100
cctcggcgag cggcccacgg tcaaccocgg tcccgggtgtt gacgcctacg tctgggtgaa 2160
gcccccggtt gagtccgacg ggcgccagcg ggagatcccg aacgacgagg gcaagggtctt 2220
cgaccgcatg tgcgaccocga cctaccaggg caacgcccgc aacggcaaca acccctcgga 2280
tgcgctgccc aacgccccca tctcoggcca ctggttctct gccagttcc gcgagctgct 2340
ggccaacgcc taccgcctc tgtaaagcgg agtgaggcaa cggctgacag cctcaacgag 2400
gaactgatca gcacctccta gccggagacg gcgcccgtcc actccccgtg ggcgggcgcc 2460
gcttttatgc cgaccctgac cccagccgca aggggcacgg gtcggcctat tccggcgatg 2520
tcggtcacgt cgccctagca cccggaaacg ccgagaaaga ctgccccgaa acggtccctct 2580
cccatccctg cattaggttg gccgagtcgg cctatggctt cgtgggcccg aacccaaccc 2640
accatcaacg agaggtatca ccatggccag tgtggtgaaa ttcaatgtgc tgacggttcc 2700
tcccggtgcc ggcgccaccc cggaggacgt ttgccaaagc gcgaggcctc gtggagaacc 2760
gggcccgggtt tcacgagttc caactgccgg cgcggcgga cgggacggac aagtacatcg 2820
tctacacgcg ctggcgctcc ggagaggact accagaactg gctgaacagc gaggccttcc 2880
agcgcggaac cgcccaggcc tctgaagact cccgcgcgag cagccagggc ggcccggccg 2940
cgtccgcgag tgaactctgg tccttcgaag tcgtccagca cgtccaggcc caggactgat 3000
cccggtgccg cctcgggttc tttaccgggg gccgcccacc cccttcatcc cttttcttct 3060
cccccgcaac ccttttgatc tgcaatgatg gaattcgca ttcttgagaa ggccgatcgt 3120
gtccatgacc ggcgagaagg caggacgacc acgcgtaccg gtcgacatcg aaggagtcaa 3180
ctgacagtgg ggactatcgc ggggctgatt gtcgcgctgt caggcgtggg gatggtctcg 3240
gccaacgtgc tcccgtggga accgtcggac ccggcatccg tgggtccccgc cacctcgag 3300
ggcagcagtt ctcccatgac gccggagccc tcgcgtcccc ggtaccccca ctcgtcgct 3360
ccgtggtcga agaggtgccc agcgcaagcg gagaactcgc ggtcgtcgaa ggtgacgggg 3420
aggctcgtcg cgaaggcacg ctctcgcgct acctggtgga ggtcgaagaa gggcttcccc 3480
gagaccccg cgaacttcgt gca
3503

```

<210> 3

<211> 21

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: VSP Leader
Sequence

<400> 3

Met Lys Leu Phe Val Phe Phe Val Ala Ala Val Val Leu Val Ala Trp

1

5

10

15

Pro Cys His Gly Ala

20

<210> 4

<211> 25

<212> DNA

<213> Artificial Sequence

;
;

<220>

<223> Description of Artificial Sequence: Xba E2 PCR
Primer

<400> 4

gctctagatg aatgattctc cgttc

25

<210> 5

<211> 17

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Xba E2 PCR
Primer

<400> 5

tgaccggcag caaaatg

17

<210> 6

<211> 25

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Xba E3 PCR
Primer

<400> 6

gctctagatg gccggctgct cggtg

25

<210> 7

<211> 23

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: RIE3 PCR
Primer

<400> 7

ggaattctta cagaggcggg tag

23

<210> 8

<211> 3004

<212> DNA

<213> Acidothermus cellulolyticus

<300>

<400> 8

```

ggatccacgt tgtacaaggt cacctgtccg tcgttctggt agagcggcgg gatggtcacc 60
cgcacgatct ctcttttgtt gatgtcgacg gtcacgtggt tacggtttgc ctcgcccgcg 120
attttcgcgc tcgggcttgc tccggctgtc ggggttcgggt tggcgtggtg tgcggagcac 180
gccgaggcga tcccaatgag ggcaagggca agagcggagc cgatggcacg tcgggtggcc 240
gatggggtac gccgatgggg cgtggcgctc ccgccgcgga cagaaccgga tgcggaatag 300
gtcacgggtgc gacatgttgc cgtaccgcgg acccgatga caaggggtggg tgcgcgggtc 360
gcctgtgagc tgccggctgg cgtctggatc atgggaacga tcccaccatt ccccgcaatc 420
gacgcgatcg ggagcagggc ggcgcgagcc ggaccgtgtg gtcgagccgg acgattcgcc 480
catacgggtgc tgcaatgccc agcgccatgt tgtcaatccg ccaaatgcag caatgcacac 540
atggacaggg attgtgactc tgagtaatga ttggattgcc ttcttgccgc ctacgcgtta 600
cgcagagtag gcgactgtat gcggtagggt ggcgctccag ccgtgggctg gacatgcctg 660
ctgcgaactc ttgacacgtc tggttgaacg cgcaatactc ccaacaccga tgggatcggt 720
cccataagtt tccgtctcac aacagaatcg gtgcgcctc atgatcaacg tgaaaggagt 780
acggggggaga acagacgggg gagaaaccaa cgggggattg gcggtgccgc gcgcattgcg 840
gcgagtgcct ggctcgcggg tgatgctgcg ggtcggcgct gtcgtcgcgg tgctggcatt 900
ggttgccgca ctcgccaacc tagccgtgcc gcggccggct cgcgcgcggg gcggcggtta 960
ttggcacacg agcggcgggg agatcctgga cgcgaacaac gtgccgttac ggatcgccgg 1020
catcaactgg tttgggttcg aaacctgcaa ttacgtcgtg cacggtctct ggtaacgcga 1080
ctaccgcagc atgctcgacc agataaagtc gctcggctac aacacaatcc ggctgccgta 1140
ctctgacgac attctcaagc cgggcaccat gccgaacagc atcaattttt accagatgaa 1200
tcaggacctg caggggtctga cgtccttgca ggtcatggac aaaatcgtcg cgtacgcggg 1260
tcagatcggc ctgcgcacat ttcttgaccg ccaccgaccg gattgcagcg ggcagtcggc 1320
gctgtggtac acgagcagcg tctcggaggc tacgtggatt tccgacctgc aagcgtggc 1380
gcagcgctac aagggaaccc cgacggtcgt cggctttgac ttgcacaacg agccgcacga 1440
cccggcctgc tggggctgcg gcgatccgag catcgactgg cgattggccg ccgagcgggc 1500
cggaaacgcc gtgctctcgg tgaatccgaa cctgctcatt ttcgtcgaag gtgtgcagag 1560
ctacaacgga gactctact ggtggggcgg caacctgcaa ggagccggcc agtaccgggt 1620
cgtgctgaac gtgcgaacc gcctggtgta ctcggcgcac gactacgcga cgagcgtcta 1680
ccgcgagacg tggttcagcg atccgacctt cccaacaac atgcccgga tctggaacaa 1740
gaactgggga tacctcttca atcagaacat tgcaccgta tggctgggcg aattcgggtac 1800
gacactgcaa tccacgaccg accagacgtg gctgaagacg ctcgccaggt acctacggcc 1860
gaccgcgcaa tacggtgcgg acagcttcca gtggaccttc tggtcctgga accccgattc 1920
cggcgacaca ggaggaattc tcaaggatga ctggcagacg gtcgacacag taaaagacgg 1980
ctatctcgcg ccgatcaagt cgtcgatttt cgatcctgtc ggcgcgctcg catcgcttag 2040
cagtcaaccg tccccgtcgg tgtcgccgtc tccgtcgccg agcccgctcg cgagtcggac 2100
gccgacgct actccgacgc cgacagccag cccgacgcca acgctgaccc ctactgctac 2160
gcccacgccc acggcaagcc cgacgcgctc accgacggca gcctccggag cccgctgcac 2220
cgcgagttac caggtcaaca gcgattgggg caatggcttc acggtaacgg tggccgtgac 2280
aaattccgga tccgtcgca ccaagacatg gacggtcagt tggacattcg gcggaaatca 2340
gacgattacc aattcgtgga atgcagcggc cagcgagaac ggtcagtcgg taacggctcg 2400
gaatatgagt tataacaacg tgattcagcc tggtcagaac accacgttcg gattccaggc 2460
gagctatacc ggaagcaacg cggcaccgac agtcgcctgc gcagcaagtt aatacgtcgg 2520
ggagccgacg ggaggggtccg gaccgtcgggt tccccggctt ccacctatgg agcgaacca 2580
acaatccgga cggaactgca ggtaccagag aggaacgaca cgaatgcccg ccatctcaaa 2640

```

;

acggctgcga gccggcggtcc tcgcccggggc ggtgagcatc gcagcctcca tcgtgccgct 2700
ggcgatgcag catcctgccca tcgcccgcgac gcacgtcgac aatccctatg cgggagcgac 2760
cttcttcgtc aaccgcgtact gggcgcaaga agtacagagc gaacggcgaa ccagaccaat 2820
gccactctcg cagcgaaaat gcgcgtcggt tccacatatt cgacggccgt ctggatggac 2880
cgcatcgctg cgatcaacgg cgtaacgggc ggaccggct tgacgacata tctggacgcc 2940
gccctctccc agcagcaggg aaccaccct gaagtcattg agattgtcat ctacgatctg 3000
ccgg 3004

<210> 9

<211> 2220

<212> DNA

<213> *Trichoderma reesei*

<300>

<308> Genbank E00389

<309> 1997-09-29

<310> JP 1985149387-A1

<312> 1985-08-06

<400> 9

aaggtagcc aagaacaata gccgataaag atagcctcat taaacggaat gagctagtag 60
gcaaagtcag cgaatgtgta tatataaagg ttcgagggtcc gtgcctccct catgctctcc 120
ccatctactc atcaactcag atcctccagg agacttgtag accatctttt gaggcacaga 180
aacccaatag tcaaccggcg actggcatca tgtatcggaa gttggccgtc atcacggcct 240
tcttgggccac agctcgtgct cagtcggcct gcactctcca atcggagact caccgcctc 300
tgacatggca gaaatgctcg tctggtggca cttgcactca acagacaggc tccgtggtca 360
tcgacgccaa ctggcgctgg actcacgcta cgaacagcag cacgaactgc tacgatggca 420
acacttgagg ctgcacccta tgcctgaca acgagacctg cgcgaagaac tgctgtctgg 480
acgggtgccgc ctacgcgtcc acgtacggag ttaccacgag cggtaacagc ctctccattg 540
gctttgtcac ccagtctgcg cagaagaacg ttggcgctcg cctttacctt atggcgagcg 600
acacgacctc ccaggaattc accctgcttg gcaacgagtt ctctttcgat gttgatgttt 660
cgcagctgcc gtaagtgaact taccatgaac ccctgacgta tcttcttggtg ggctccagc 720
tgactggcca atttaagggtg cggcttgaac ggagctctct acttcgtgtc catggacgog 780
gatggtggcg tgagcaagta tcccaccaac aacgtggcg ccaagtacgg cacggggtag 840
tgtgacagcc agtgtccccg cgatctgaag ttcacatg gccaggccaa cgttgagggc 900
tgggagccgt catccaacaa cgcaaacacg ggcattggag gacacggaag ctgctgctct 960
gagatggata tctgggaggc caactccatc tccgaggctc ttacccccca ccttgcaag 1020
actgtcggcc aggagatctg cgagggtgat gggcgcgcg gaacttactc cgataacaga 1080
tatggcgcca cttgcgatcc cgatggctgc gactggaacc cataccgcct gggcaacacc 1140
agctttctacg gccctggctc aagctttacc ctcgatacca ccaagaaatt gaccgttgct 1200
accagttcg agaagtcggg tgccatcaac cgatactatg tccagaatgg cgtcactttc 1260
cagcagccca acgcccagct tggtagttac tctggcaacg agctcaacga tgattactgc 1320
acagctgagg agacagaatt cggcggtatc ctttctcaga caaggcggc ctgactcagt 1380
tcaagaaggc tacctctggc ggcatggttc tggatcatgag tctgtgggat gatgtgagtt 1440
tgatggacaa acatgcgcgt tgacaaagag tcaagcagct gactgagatg ttacagtact 1500
acgccaacat gctgtggctg gactccacct acccgacaaa cgagacctcc tccacaccgc 1560
gtgcgctgcg cggaagctgc tccaccagct ccggtgtccc tgctcaggtc gaatctcagt 1620
ctcccaacgc caaggtcacc ttctccaaca tcaagttcgg acccattggc agcaccggca 1680

```

accctagcgg cggaaccct ccggcgga accgtggcac caccaccacc cgccgcccag 1740
ccactaccac tggaagctct ccggaccta ccagctctca ctacggccag tgcggcggtg 1800
ttggctacag cgccccacg gtctgcgcca gcggcacaac ttgccaggtc ctgaaccctt 1860
actactctca gtgcctgtaa agctccgtgc gaaagcctga cgcaccggtg gattcttggg 1920
gagcccgat catgacggcg gcgggagcta catggccccg ggtgatttat tttttttgta 1980
tctactcttg acccttttca aatatacggc caactcatct ttcactggag atgcggcctg 2040
cttggtattg cgatgttgtc agcttgga attgtggctt tcgaaaacac aaaacgattc 2100
cttagtagcc atgcatttta agataacgga atagaagaaa gaggaatta aaaaaaaaaa 2160
aaaaacaaac atcccgttca taaccgtag aatcgccgct cttcgtgtat ccagtagcca 2220

```

<210> 10

<211> 21

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Nar El PCR
Primer

<400> 10

cggggcgcgc gcggcggcta t 21

<210> 11

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Sac El PCR
Primer

<400> 11

cagagctctt aacttgctgc 20

<210> 12

<211> 25

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Sac Elcd PCR
Primer

<400> 12

tggagctcta gacaggatcg aaaat 25

<210> 13

<211> 76

<212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: VSP leader
 coding sequence

<400> 13
 tctagagtcg accatgaagt tgtttgtttt ctttggtgct gcagtagttt tggtagcatg 60
 gccatgccat ggcgcc 76

<210> 14
 <211> 24
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: CBH2-2 PCR
 Primer

<400> 14
 gctctagatg tatcggaagt tggc 24

<210> 15
 <211> 24
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: CBH3-1 PCR
 Primer

<400> 15
 cccccggggtt acaggcactg agag 24

<210> 16
 <211> 2199
 <212> DNA
 <213> Cellulomonas fimi

<300>
 <303> Gene
 <304> 44
 <306> 315-342
 <307> 1986
 <308> Genbank M15823
 <309> 1993-06-04

<400> 17
ggctgcaggc ggctgccgcg tcgac 25

<210> 18
<211> 23
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: cemAsac PCR
Primer

<400> 18
ccgagctctc accacctggc gtt 23

<210> 19
<211> 82
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: VSP leader
coding sequence

<400> 19
tctagagtgc accatgaagt tgtttgtttt ctttgttgca gcagtagttt tggtagcttg 60
gccttgccac ggcgctgcag tc 82

<210> 20
<211> 2286
<212> DNA
<213> Clostridium thermocellum

<300>
<303> Nucleic Acids Res.
<304> 14
<305> 21
<306> 8605-8613
<307> 1986
<308> Genbank X04584
<309> 1999-02-10

<400> 20
aaactaaaac tcctatccaa tacttttagtt cagttccagc atacgtctgt attcaaaatg 60
cotgtatttta taactgcatt tataatacct gaagcaaata ataattaaac ttgtggaaga 120
aaggaggttg caacagggttt taaattatct taattcaggt attttacaat ttttaataaa 180
aaggggggata aaggtaaaaa atgagtagaa tgaccttgaa aagcagcatg aaaaaacgtg 240
tgttatcttt gctcattgct gtagtgtttc taagcttgac cggagtattt ccttcgggat 300

```

tgattgagac caaagtgtca gctgcaaaaa taacggagaa ttatcaattt gattcacgaa 360
tccgtttaaa ctcaataggt tttataccga accacagcaa aaaggcgact atagctgcaa 420
attggtcaac cttttatggt gttaaagaag acggaacaat agtgtatacc ggaacggcaa 480
cttcaatggt tgacaatgat acaaaagaaa ctgtttatat tgctgatttt tcatctgtta 540
atgaagaagg aacgtactat cttgccgtgc cgggagtagg aaaaagcgta aactttaaaa 600
ttgcaatgaa tgtatatgag gatgctttta aaacagcaat gctgggaatg tatttgctgc 660
gctgcggcac cagtgtgtcg gccacataca acggaatata ctattcccat ggaccgtgcc 720
ataactaatga tgcatactct gattatataa acggacagca tactaaaaaa gacagtacaa 780
aaggctggca tgatgcgggc gactacaaca aatatgtggt aaacgccggc ataaccgttg 840
gttcaatggt cctggcgtgg gagcatttta aagaccagtt ggagcctgtg gcattggaga 900
ttcccgaaaa gaacaattca ataccggatt ttcttgatga attaaaatat gagatagact 960
ggattcttac catgcaatac cctgacggga gcggaagggt ggctcataaa gtttcgacaa 1020
ggaacttttg cggctttatc atgcctgaga acgaacacga cgaaagattt ttctgtccct 1080
ggagcagtg cgcacacggca gactttgttg ccatgacggc catggctgca agaataattca 1140
ggccttatga tcctcaatat gctgaaaaat gtataaatgc ggcaaaagta agctatgagt 1200
ttttgaagaa caatcctgcg aatgtttttg caaaccagag tggattctca acaggagaat 1260
atgccactgt cagtgatgca gatgacagat tgtgggcggc ggctgaaatg tgggagaccc 1320
tgggagatga agaatacctt agagattttg aaaacagggc ggcgcaattc tcgaaaaaaa 1380
tagaagccga ttttgactgg gataatgttg caaacttagg tatgtttaca tatcttttgt 1440
cagaaagacc gggcaagaat cctgcttttg tgcagtcaat aaaggatagt ctcctttcca 1500
ctgcggatcc aattgtgagg accagccaaa accatggcta tggcagaacc cttggtacaa 1560
catattactg gggatgcaac ggcacggttg taagacagac tatgatactt caggttgcga 1620
acaagatttc acccaacaat gattatgtaa atgctgctct cgatgcgatt tcacatgtat 1680
ttggaagaaa ctattacaac aggtcttatg taacaggcct tggataaat cctcctatga 1740
atcctcatga cagacgttca ggggctgacg gaatatggga gccgtggccc ggttaccttg 1800
taggaggagg atggcccgga ccgaaggatt gggtgatat tcaggacagt tatcagacca 1860
atgaaattgc tataaactgg aatgcggcat tgatttatgc ccttgccgga tttgtcaact 1920
ataattctcc tcaaaatgaa gtactgtacg gagatgtgaa tgatgacgga aaagtaaact 1980
ccactgactt gactttgtta aaaagatatg ttcttaaagc cgtctcaact ctcccttctt 2040
ccaaagctga aaagaacgca gatgtaaata gtgacggaag agttaattcc agtgatgtca 2100
caatactttc aagatatttg ataagggtaa tcgagaaatt accaatataa attctgataa 2160
atattgataa acactaatat ataagtgtt aatcggtaaa agagccctgt ggcaaaaact 2220
gcgcagaggt gtttttatca attccggcgc agacgaaaat agcagacgta aatattaatt 2280
actgaa
2286

```

<210> 21

<211> 17

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: celDpst PCR
Primer

<400> 21

agctgcagaa ataacgg

17

<210> 22

<211> 26
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: celDsac PCR
 Primer

<400> 22
 ccgagctctt atattggttaa tttctc

26

<210> 23
 <211> 4578
 <212> DNA
 <213> Clostridium cellulovorans

<300>
 <303> Gene
 <304> 211
 <305> 1
 <306> 39-47
 <307> 1998
 <308> Genbank U34793
 <309> 1999-03-09

<400> 23
 taatatataa ataatatatta aaatcaataa atcaatcggg gaaaatttaa aaaaagagag 60
 ggataatcaa tgagaaaaag attaaataag atcgttgctg ttgctttaac tgcaacaact 120
 atatcatcag tagcagctac tgtaataaca gctcaagttt cagctgcacc agtagtgcca 180
 aataatgagt atgttcaaca ctttaaggat atgtacgcta agatccataa tgcaacaact 240
 ggataacttca gtgatgaagg aataccttat cacgcagttg aaacattaat gggtgaagca 300
 ccagactatg gtcatgaaac tacaagtga gcttgggatg taactgaaaa gtacataatt 360
 ccaggtgaga ctgatcaacc aagcgcaagt atgagcaatt atgatccaaa taagccagct 420
 acatatgcag ctgaacatcc agatccaagc atgtacccat ctcaattaca atttggtgct 480
 gctgtaggta aggatccatt atacaatgaa ttaaaatcta cttatggaac tagccaagta 540
 tatggtatgc attgggttact agatgttgat aactggtatg gttttggtgg tgcaacaagc 600
 acaagcccag tatacatcaa cactttccaa agaggtgttc aagaatcttg ttgggaaact 660
 gtgccacaac catgtaaaga cgaaatgaag tacggtggaa gaaacggttt cttagatcta 720
 ttcaactggtg attcacaata cgcaactcaa tttaaataata ctaacgctcc agacgcagat 780
 gctcgtgcag ttcaagctac ttactatgca caattagctg ctaaagaatg gggagtagac 840
 atcagctcat atgtagcaaa atctactaag atgggtgact tottaagata ttcattcttt 900
 gataaatact ttagaaaagt tggaaattca acacaagcag gaactggata tgattcagct 960
 caatacctat taaactggta ctatgcttgg ggtggtggaa tcagctcaaa ctggtcttgg 1020
 agaattggat caagccataa ccatttcgga taccaaaacc caatggcagc atggatatta 1080
 tcaaatacat ctgactttta accaaagtca ccaaagctg ctacagattg gaataacagt 1140
 ttaaagagac aaatagaatt ctatcaatgg ttacaatctg ctgaaggtgg tatcgttgga 1200
 ggagctagta actcaaagtg aggaagctat caagcatggc cagcaggtac tgaacattc 1260
 tacggaatgg gatatactcc tcaccagta tacgaagatc caggtagtaa cgaatggttt 1320


```

ggtatgcaag catggtcaat gcaacgtgtg gctgaatact actacagttc aaaagatcca 1380
gcagctaaat cattacttga taaatgggct aaatgggctt gtgcaaagt tcaattcgat 1440
gatgcagcta agaaatttaa gattcctgct aaattagtat ggactggaca accagatact 1500
tggactggat catatacagg aaattcaaat cttcatgtta aagttgaagc ttatggagaa 1560
gatcttggag tagcagggttc acttttcta gctattatcat attatgcaaa agctcttgaa 1620
tctagcacag atgctgcaga taaagtagca tataacactg caaaagaaac ttctagaaaag 1680
atacttgatt acttatgggc aagctaccaa gatgataagg gtatagcagt tactgaaaca 1740
agaaatgatt tcaaacgttt caatcaatct gtatatattc catcaggttg gacaggaaaa 1800
atgcctaata gagatgtaat ccaaagtgga gctactttct taagcatacg ttcaaaatac 1860
aaacaagatc catcatggcc aaatggtgaa gctgctttag caaatggtac tgggtgttgat 1920
atgacatacc acagattctg gggctcaaagt gatatcgcta tagcatttgg aacatacggg 1980
acattattca cagaccctac tccaggatta aaagggtgatg ttaactctga tgctaaagta 2040
aatgctatag attttagctat attaaagaaa tacatcttag attcaacaac taaaattaac 2100
actgctaatt ctgatatgaa cgggtgatgga aaagttaatg caatggattt agctttatta 2160
aagaaagcac ttctgcttaa gattaaataa ctttagatcg aaattgtaag gttattttaag 2220
gctggacaat atcaagtata ttgtccagct actttaaaaa atattgggaa acactgtgta 2280
aggtaaaact aaaccatgga tatgaaatat agtaagatta atgccattgc tatggcaaac 2340
ttaaataaaa tatattagag cataaacatg aaatttaagt aaaaggcgaa taaataattc 2400
cctaatacaa aaaattaagg ggtggaacta gtgtttaaca tatctaagaa aaaagcgcaa 2460
gctcttcttt tatcaggaat cttgggtgca acttcattta caccagctgt attggtaaaa 2520
ggtgaaacaa cagcgactcc aacattcaat tatggagaag cattacaaaa gtcaataatg 2580
ttttatgaat tccaacgttc tggaaagtta ccaacggata ttcgtagtaa ttggcgtggg 2640
gattctggaa caaagatgg ctctgatgta ggagttgatt taactgggtg atggtatgat 2700
gctggagacc acgttaaatt taatctgcca atgtcttata ctgtggcaat gcttgcattg 2760
tcattaagtg aagacaaagc agcttacgaa aaaagcgcc aattagatta ccttggttaag 2820
gaaataaaat gggctacaga ttatctaatt aagtgccata cggcaccaaa tgaatactat 2880
tatcaagttg gtgatggtgg agctgatcac aaatggtggg gacctgcaga agtaatgcag 2940
atggcaagac cggcttataa agtagatttg caaaaaccag gatcatcagt tgctgctgaa 3000
acagcagcag cattagcttc tacagctttt gcattaaaag acatagataa agcgtattca 3060
gaacaatgta ttcagcatgc aaaagaactt tataactttg ctgatacaac aaagagtgat 3120
gctggttata cagcagcaaa tacatattac aattcatgga gtggatacta tgatgaatta 3180
tcatgggctg cagcatggct ttacatggca acaaatgatg catcatatct agaaaaagcg 3240
gaatcatatg ttccattttg gaaggttgaa cagcaaacaa ccactatagc atatagatgg 3300
gogcattgtt gggatgatgt acatttcgga gctcaattac tccttgccag attaacagga 3360
aaatcaatat acaaagaatc agttgaaaga aaccttgatt attggacaac tggttatgat 3420
ggaaataaaa taaagtacac tccaaaagg ttagcttgga tggattcttg gggctcatta 3480
agatatgcaa ctacaacggc attccttgcc gatgtttatg caagctcaga tgtttgttct 3540
atctctaagg tagatacata taagaatttt gctaagagtc aagctgatta tgcttttagg 3600
agtactggaa gaagttttgt ggtaggattt ggtgaaaatg ctccaaagaa accacatcat 3660
agaactgcc atagttcatg gtcagatcaa caagtaaata caacagacca tagacatgtt 3720
ttatatggtg ctttagttgg aggaccagat gccagtgatg gttatactga tgctattgac 3780
aattttacta ataatgaggt ggcttgatg tataatgcag gatttgtagg acttttagct 3840
agacaatat ctaaatatgg cggagatcca atacctgatt ttaaagcgat agaaaagcca 3900
accaacgatg agttctttgt cgaagcagga gtaaatgta cagggtccaaa ttttgtagaa 3960
attaaagctt tagttaataa tagaacagga tggccagcaa gaatgggaga taaactttca 4020
ttcaataact tcataaatgt aagtgaattt gttaatgctg gttacagtgc agatgattta 4080
aaggttactg ttggttacia tactggcgga actgtatcaa acctaatccc atgggataag 4140
gaaaataata tttattatgt aaatgttgat ttcacagggg taaagattta tccagggtgga 4200

```

```

caatcagatt ataaaaaaga aattcaattt agaatttcag gaattcaaaa tgttaatatt 4260
tgggataatt ctgatgactt ctcttatgag gggattacaa aaactccagg tgaaacacct 4320
gtgaagggtta caaacatccc agtttatgat aatggagtta aggtattcgg aaatgaacca 4380
ggaactacta agccacctgt tatagctggg gatgtaaaca ataatgggtat cgtgaattca 4440
atggatttag cgatgtttaa gaaatatata ctgggatacg aagtagaaat gaataaagag 4500
gcttcagatt taaataaaga tggttaagatt aatgccattg atttcgctct tttaaagaaa 4560
ctacttttat cacagtag 4578

```

<210> 24

<211> 24

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: exgSnar PCR
Primer

<400> 24

cggggcgccg caccagtagt gcca 24

<210> 25

<211> 24

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: exgSsac PCR
Primer

<400> 25

cagagctctt atttaatctt aagc 24

<210> 26

<211> 5971

<212> DNA

<213> Thermobifida fusca

<300>

<400> 26

```

gcgggcgcca ggtggggcgg ccgagtcgga cggtggcgag gacttcgacc ccgtgggtcgt 60
cagcgaagca tgcttcgctg acgttggcgg cggtagcag gcctaccccg gtgccgggtca 120
tcccgaagtg ggtggcgagt tcttcgaggt ggttggtccgg gtccggtccgc cggtagtcgc 180
cgtcgacttg ggcgttgatg acgtagtccc gtggtccgag gccgccgcgc agcacgctgg 240
aggcgggccat gcgccagccg gggccgcact gccagagcac tgcggcaagg gcgcgggccgt 300
tgtcgcgcgc ccagcgcagg tgcgcggtga ggagggtgcc gtccgcgata gtctccgtgt 360
ccactccgct ctccagggtc ggttcgcgcg ggccacttta acgagtcgc atcggggcggc 420
agcgggggtca tgcttccatc tgggcgaatg ccacgcggag tactgcgagt ctcacgcttc 480

```

```

gtccattgt ctttttacgg aggggtttaca cggcacatcc ggaacgttac cctcctactg 540
ggagcgctcc cgtgactcag gccacaggcc cccttcccgg ggcgagcccc caccaccccg 600
ggaaggccac tgcgaccctt gagctcaatt catgggagcg ctcccatgcc agtgctcccg 660
gcagaaagga gagaaccgga acccggatcc gctggtttca ctgatcccc cgtttttacc 720
cggcatcgcc ccgacaccag cgatgccacc cccgcggcgg ctgcagtcca cacccggcac 780
gocgcacacc cctctcctgt gtgacacgcg gttccgaacg gccttgccgg ttctcgggat 840
gccgacgaac ggcacggctg ctccggaacc ggaagatccg gaggttattt ccaagcatga 900
gatcgttact gtctccccgg cgctggcgca cgctggcctc gggggcgctc gcagcgggcc 960
tgcccgccgc tgtactctcc cccggcgctg cgcacgccgc cgtcgccctg tgggtggact 1020
acgacgactc caacgactgg ggtagcgggt tcgtcgccga agtcaagggt accaacgaag 1080
gcagcgaccc catccagaac tggcaagtag gctggacctt ccccggtaac cagcagatca 1140
ccaacggctg gaacggcggt ttccagccaga gcggcgccaa cgtcacccgt cgtaccggg 1200
actggaaccc caatatcgcc cccggagcca ccatctcctt cggcttccag ggcacctaca 1260
gocgctccaa cgacgccccg accagcttca ccgtcaacgg cgtcacctgc agcggatccc 1320
agcccgccaa cctgcgcgcc gatgtcacc tgacatcccc ggccaacaac tcgaccttcc 1380
tggtcaacga cccgatcgag ctgaccgcgg tcgcctccga ccccgacggc tcgatcgacc 1440
gggtggaatt cgcgcgcgac aacaccgtca tcggcatcga caccacctcc cctacagct 1500
tcacctggac ggacgctgcc gccggctcct actcggtgac cgcgatcgcc tacgacgacc 1560
agggagccag gaccgtctcc gctcccatcg ccatccgagt gctggaccgg gccgcgctca 1620
tcgectcacc gccacccgtc cgcgtgccgc agggcggcac cgcgcacttc gaggtgcggc 1680
tgtccaacca gccctccggc aacgtcacgg tcacctggc gcgcacgtcg ggcagctccg 1740
acctgaccgt ctccagcggc tcccaactcc agttcacctc cagcaactgg aaccagccgc 1800
agaaggtgac catgcgctcc gctgacaacg gcggaacct ggccgaggcg gtcttcaccg 1860
tcagcgcccc cggccacgac tcggccgagg tgacggtcgg ggagatcgac ccgaacacca 1920
gtcctacga ccaggccttc ctggagcagt acgagaagat caaggacccc gccagcggct 1980
acttccgcga attcaacggg ctcttggtcc cctaccactc ggtggagacc atgatcgctg 2040
aggctccgga ccacggccac cagaccagt ccgaggcgtt cagctactac ctgtggtctg 2100
aggcgtaacta cggccgggtc accggtgact ggaagccgct ccacgacgcc tgggagtcca 2160
tgagagacct catcatcccc ggcaccaagg accagccgac caactccgcc tacaaccca 2220
actccccggc gacctacatc cccgagcagc ccaacgctga cggctacccg tcgctctctc 2280
tgaacaacgt cccggtgggt caagaccgc tcgcccagga gctgagctcc acctacggga 2340
ccaacgagat ctacggcatg cactggctgc tcgacgtgga caacgtctac ggcttcgggt 2400
tctgcggcga cggcaccgac gacgcccccg cctacatcaa cacctaccag cgtggtgcgc 2460
gcgagtcggg gtgggagacc attccgcacc cgtcctgcga cgacttcacg cagggcggcc 2520
ccaacggcta cctggacctg ttaccgacg accagaacta cgccaagcag tggcgctaca 2580
ccaacgcccc cgacgctgac gcgcggggcg tccagtgat gttctggggc cacgaatggg 2640
ccaaggagca gggcaaggag aacgagatcg cgggctgat ggacaaggcg tccaagatgg 2700
gcgactacct ccggtacgag atgttcgaca agtacttcaa gaagatcggc aactgcgtcg 2760
gcgcacacct ctgcccgggt ggccaaggca aggacagcgc gcactacctg ctgtcctgg 2820
actactcctg gggcggtcg ctcgacacct cctctgcgtg ggcgtggcgt atcggtcca 2880
gtcctcgca ccagggttac cagaacgtgc tcgctgccta cgcgtctctg caggtgcccg 2940
aactgcagcc tgactccccg accggtgtcc aggactgggc caccagcttc gaccgccagt 3000
tgaggttctt ccagtggctg cagtccgctg aagtggttat cgcgggtggc gccaccaaca 3060
gctggaaggg aagctacgac accccgccga ccggcctgtc gcagttctac ggcatgtact 3120
acgactggca gccggtctgg aacgaccgc cgtccaacaa ctggttcggc ttccaggtct 3180
ggaacatgga gcgcgtcgcc cagctctact acgtgaccgg cgacgcccgg gccgaggcca 3240
tctctgacaa gtgggtgccg tgggccatcc agcacaccga cgtggacgcc gacaacggcg 3300
gccagaactt ccagggtccc tccgacctgg agtggtcggg ccagcctgac acctggaccg 3360

```

```

gcacctacac cggcaacccg aacctgcacg tccaggctcg ctccctacagc caggacgtcg 3420
gtgtgaccgc cgctctggcc aagaccctga tgtactacgc gaagcggttcg ggcgacacca 3480
ccgccctcgc caccgcggag ggtctgctgg acgccctgct ggcccacccg gacagcatcg 3540
gtatcgccac ccccgagcag ccgagctggg accgtctgga cgaccctggg gacggctccg 3600
agggcctgta cgtgccgcgc ggctggctcg gcacatgcc caacgggtgac cgcctcgagc 3660
cgggcgcgac ctctctgtcc atccgctcgt tctacaagaa cgaccctgctg tggccgcagg 3720
tcgaggcaca cctgaacgac ccgcagaacg tcccggcgcc gatcgtggag cgccaccgct 3780
tctgggctca ggtggaaatc gcgaccgctg tcgcagccca cgacgaactg ttcggggccg 3840
gagctccctg atcctgaaca gtccgtctcc tgaccggcgg tccgtggcgg cgtagtgtctc 3900
ccccgcgcgc gaccgcctc caccacccc ctggggccgt ccggcgcacc aacaccggac 3960
ggccccagac tcttttcacg gctgtctact cggcccgcag ccccgcccta tccagcccga 4020
tcaaccaggt cagcgtagtc cgggtgccgc tcgagccacg ccttgacata ggggcagacc 4080
gcacgcaccc gcagccgcgc ggccgcgaca tcatccaacg cctcccgcac cagccggctg 4140
cccatccctc tcccctggaa agagccgtcc acccgagtgt ggaagaagac caccgatca 4200
ccggcaacag gctggtatcc cgtgaaaccg gcgattttct ccccggaacg gatctcgta 4260
cgggtgctgtg ccgattgtc gatcaccgtg acatccatgg cgttccctc cgttcccca 4320
ggcggcgcgc catctctcat ccggcggcct gccaggact gtctctctcc ccaggacagc 4380
ctgagcacgc gccgcacgg cgaaccccag ccacagccct agcatggtga acggccacgg 4440
tcgcagggcg accacgacac cgaggaggag tgatgggcga cggcggcaca gtacgccacc 4500
cttccgtacg tgccgtatca ccggcagcgt tggacgaggc ggtaacgcgtc ctgcgggcag 4560
gcggcctcgt agctttcccc accgaaaccg tctacggcct aggcgcggac gccgcgaacc 4620
cggcgcgcgt agcgcgcatt ttgcgcgcca aaggccgtcc cgctgaccac cccctgatcg 4680
tgacgtcgc ttccgcggag agcgcgcgcg actgggctgc tacctttcct cctctcgccc 4740
gcacgtggc ggacgcgttc tggcccggcc cgctgaccct cactctcccc cgctccgccc 4800
aagtgcgcga cgcagtgacc ggtggacgcg ccacggtagg gctgcgggtc cccgaccagc 4860
cggtcgcgct cgcactcctg gaacgcttcg gcgcgcgaat cgcgcgaact tccgcgaacc 4920
ggttcggccg agtgagcccc accacggccg cgcacgttgc cgctgacctc ggggaccggg 4980
tcgacctggt gttggacggc ggaccgtgca cggtcggcgt ggaatcgacg atcgtcgaag 5040
tggccgacgg ccggctcacc gtgctccgca ccggaggcat caccoccgac gaccttgccg 5100
cggtcaccgg agccccgcgc gacaccacc ccaccggacc ggcccgggca cccggcatgc 5160
tcgcccgcga ctacgcaccc gccgcacggg tcgtgctagc cgaagcagca gaagccgcgg 5220
acacggtcgc ccagtggttg gagaaaggac accgggtggc tgtgctggcg gagaccgcta 5280
ccgtgcccga aaacctgccg gaaggcgtgg tgggtctacc gtcccctgct tcggctcggg 5340
actatgcccg cgtgctgtac cagcggtgcg gggacgtgga cgcggcggga gccgacgtgg 5400
tcgtcgcgat ccccccgaa cccgcgggga tcggcttggc ggtacgggac cggctgctgc 5460
gcgcatcccg ggcgactga cctctcccc tggggcaagg gatTTTTccg catagacgag 5520
cccgTTTTcc ggatctcttt ctgtggagac agaaagagag caccgacacc agggaggggc 5580
gatgaaagct caagccggag accggatcgt tgtggaacgc ccccgcgatg acctgcccgc 5640
gcgcaaaggc gtcgtgctca aagtgcaggg ggacaacggg ggcccgcct actgggtgcg 5700
gtgggaggac gaaggccggg aaacctcgt ctaccgggga ccggacgccc gtatcgagcc 5760
ccgccacccc gtcccccaag cccgccagga gcatacggag gcccgtaac cgcagccggg 5820
acagtccctg aaacggatac agatcgacgt cgcagtgtcg gaggtgcacg agaacggctc 5880
cgtgcgcacc ctgcgcgaag cgcaactgcc gtccaccaag tggaacctgc gcggccacgg 5940
agaagcacgc aagcatccca ccgacgcgca t
5971

```

<210> 27
 <211> 21
 <212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: celFpst PCR
Primer

<400> 27

acgctgcagt cgctgctcg g

21

<210> 28

<211> 23

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: celFhma PCR
Primer

<400> 28

cccccggttc agggagctcc ggc

23

<210> 29

<211> 15

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: 2777 PCR
Primer

<400> 29

ggccacctgg gcagg

15

<210> 30

<211> 17

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: M13-20 PCR
Primer

<400> 30

gtaaaacgac ggccagt

17

<210> 31

<211> 15

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: 3227 PCR
Primer

<400> 31

gcgacgctcg ggccg

15

<210> 32

<211> 16

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Reverse 3227
PCR Primer

<400> 32

aacagctatg accatg

16

DECLARATION AND POWER OF ATTORNEY

I the undersigned inventor(s), hereby declare(s) that:

My residence, post office address and citizenship are as stated below next to my name.

I believe that I am the original, first, and joint (sole) inventor(s) of the subject matter which is claimed and for which a patent is sought on the invention entitled:

TRANSGENIC PLANTS AS AN ALTERNATIVE SOURCE OF LIGNOCELLULOSIC-DEGRADING ENZYMES

the specification of which is attached hereto.

I hereby state that I have reviewed and understand the contents of the above-identified specification, including the claims, as amended by any amendments referred to above.

I acknowledge the duty to disclose information material to the examination of this application as defined in Section 1.56 of Title 37 Code of Federal Regulations.

I hereby claim the benefit under Section 120 or Section 119 (e)(1) of Title 35 United States Code, of any United States application(s) listed below and, insofar as the subject matter of each of the claims of this application is not disclosed in the prior United States application in the manner provided by the first paragraph of Section 112 of Title 35 United States Code. I acknowledge the duty to disclose material information as defined in Section 1.56(a) Title 37 Code of Federal Regulations, which occurred between the filing date of the prior application and the national or PCT international filing date of this application:

Application Serial No.	Filing Date	Status (pending, patented abandoned)
<u>08/883,495</u>	<u>June 26, 1997</u>	<u>Pending</u>

And I hereby appoint:

Charles S. Sara	30,492
Joseph T. Leone	37,170
Craig A. Fieschko	39,668
Mary E. Eberle	43,599

Address all telephone calls to: **Joseph T. Leone**
Telephone: 608-831-2100
Facsimile: 608-831-2106

All correspondence to:

Intellectual Property Department
DEWITT ROSS & STEVENS, S.C.
Firststar Financial Centre
8000 Excelsior Drive Suite 401
Madison, WI 53717-1914

I declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code and that such willful false statements may jeopardize the validity of the application or any patent issued thereon.

FULL NAME OF SOLE OR FIRST INVENTOR Sandra Austin-Phillips

INVENTOR'S SIGNATURE _____ DATE _____

Residence: Madison, Wisconsin

Citizenship: Great Britain Post Office Address: 2137 Fox Avenue, Madison, WI 53711

FULL NAME OF JOINT OR SECOND INVENTOR Richard R. Burgess

INVENTOR'S SIGNATURE _____ DATE _____

Residence: Madison, Wisconsin

Citizenship: U.S.A. Post Office Address: 10 Knollwood Court, Madison, WI 53713

FULL NAME OF JOINT OR THIRD INVENTOR Thomas L. German

INVENTOR'S SIGNATURE _____ DATE _____

Residence: Hollandale, Wisconsin

Citizenship: U.S.A. Post Office Address: 1671 Sandy Rock Road, Hollandale, WI 53544

FULL NAME OF JOINT OR FOURTH INVENTOR Thomas Ziegelhoffer

INVENTOR'S SIGNATURE _____ DATE _____

Residence: Madison, Wisconsin

Citizenship: U.S.A. Post Office Address: 418 Oak Crest Avenue, Madison, WI 53705

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

Appln. Serial No.:

Attorney Docket No.: 09820.114

Filing Date: Simultaneously Herewith

Applicants: AUSTIN-PHILLIPS et al.

Title: **TRANSGENIC PLANTS AS AN ALTERNATIVE SOURCE OF
LIGNOCELLULOSIC-DEGRADING ENZYMES**

**SUBMISSION OF "SEQUENCE LISTING," COMPUTER READABLE COPY,
AND/OR AMENDMENT PERTAINING TO A BIOTECHNOLOGY INVENTION
CONTAINING A NUCLEOTIDE AND/OR AMINO ACID SEQUENCE**

Box: Sequence

Assistant Commissioner for Patents
Washington, D.C. 20231

To the Commissioner:

Enclosed are:


1. A paper copy of the "Sequence Listing" for the nucleotide sequence in this application. The "Sequence Listing" is assigned a separate identifier as required in 37 C.F.R. 1.821(c) and 37 C.F.R. 1.822 and 1.823.
2. A copy of the "Sequence Listing" in computer readable form in accordance with the requirements of 37 C.F.R. 1.821(e) and 1.824.
3. A statement that the content of the paper copy of the "Sequence Listing" and the "Sequence Listing" in computer readable form are the same as required by 37 C.F.R. 1.821(g).

**STATEMENT THAT "SEQUENCE LISTING" AND COMPUTER READABLE COPY
ARE THE SAME AND THAT PAPERS SUBMITTED INCLUDE NO NEW MATTER**

The paper copy of the "Sequence Listing" submitted in this application is identical to the computer readable copy of the "Sequence Listing" being submitted herewith.

Should the Examiner have any questions or comments with respect to the above-referenced application, the Examiner is requested to contact the undersigned attorney. The Commissioner is authorized to charge any fees or credit any overpayments relating to this application to deposit account number 18-2055.

Respectfully submitted,



Joseph T. Leone, Reg. No. 37,170
DEWITT ROSS & STEVENS, S.C.
Firststar Financial Centre
8000 Excelsior Drive, Suite 401
Madison, Wisconsin 53717-1914
Telephone: (608) 831-2100
Facsimile: (608) 831-2106

I hereby certify that this correspondence is being deposited with the U.S. Postal Service as first class mail in an envelope addressed to:

Box: Sequence
Assistant Commissioner for Patents
Washington, D.C. 20231

Date of Deposit: 8-12-99

Signature: Maura Layton